

🔇 Tel: 400-999-8863 💌 Emall:Upingbio.163.com

Website: www.upingBio.com

TOB1 (phospho Ser164) Polyclonal Antibody

BTG family.,subunit:Interacts with CNOT7.,tissue specificity:Ubiquitous.,BackgroundThis gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have		
Reactivity Human;Mouse;Rat Applications WB;IHC;IF;ELISA Gene Name TOB1 Protein Name Protein Tob1 Immunogen The antiserum was produced against synthesized peptide derived from human TOB1 around the phosphorylation site of Ser164. AA range:130-179 Specificity Phospho-TOB1 (S164) Polyclonal Antibody detects endogenous levels of TOB1 protein only when phosphorylated at S164. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/S00 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm. Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic	Catalog No	YP-Ab-00215
Applications WB;IHC;IF;ELISA Gene Name TOB1 Protein Name Protein Tob1 Immunogen The antiserum was produced against synthesized peptide derived from human TOB1 around the phosphorylation site of Ser164. AA range:130-179 Specificity Phospho-TOB1 (S164) Polycional Antibody detects endogenous levels of TOB1 protein only when phosphorylated at S164. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polycional, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasm except in BTG family. Justically and/or functionally interacts with the erbB-2 receptor tyrosine Kinase. May physically and/or functionally interact with protein-tyrosine Kinase enception spically and/or functionally interact with the erbB-12 (Bocel translocation gene protein family. Justically and/or functionally interact with the erbB-	Isotype	lgG
Gene Name TOB1 Protein Name Protein Tob1 Immunogen The antiserum was produced against synthesized peptide derived from human TOB1 around the phosphorylation site of Ser164. AA range:130-179 Specificity Phospho-TOB1 (S164) Polyclonal Antibody detects endogenous levels of TOB1 protein only when phosphorylated at S164. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic Tissue Specificity Ubiquitous. Function function.Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase traception; .pti/Phosphorylated on Ser and Thr residues.;similarity:Belongs to the BTG family. Justically and/or functionally interact with protein-tyrosine kinase traception; .ptin	Reactivity	Human;Mouse;Rat
Protein Name Protein Tob1 Immunogen The antiserum was produced against synthesized peptide derived from human TOB1 around the phosphorylation site of Ser164. AA range:130-179 Specificity Phospho-TOB1 (S164) Polyclonal Antibody detects endogenous levels of TOB1 protein only when phosphorylated at S164. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic Function function:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase receptorsPTM:Phosphorylated on Ser and Thr residuessimilarity:Belongs to the BTG familysubuit:Interacts with CNOT7, tissue specificity.Ubiquitous. Background This gene encodes a member of the transducer of erbB-2.18-cell translocation gene protein family. Members of thi	Applications	WB;IHC;IF;ELISA
Immunogen The antiserum was produced against synthesized peptide derived from human TOB1 around the phosphorylation site of Ser164. AA range:130-179 Specificity Phospho-TOB1 (S164) Polyclonal Antibody detects endogenous levels of TOB1 protein only when phosphorylated at S164. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm. Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic Tissue Specificity Ubiquitous. Function function:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase receptors., PTM:Phosphorylated on Ser and Thr residues, similarity:Belongs to the BTG family, subunt:Interacts with CNOT7., tissue specific. Background This gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family a	Gene Name	TOB1
TOB1 around the phosphorylation site of Ser164. AA range:130-179 Specificity Phospho-TOB1 (S164) Polyclonal Antibody detects endogenous levels of TOB1 protein only when phosphorylated at S164. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic Tissue Specificity Ubiquitous. Function function: Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase receptorsPfM:Phosphorylated on Ser and Thr residues, similarity: Belongs to the BTG family, submit: Interacts with CNOT7, tissue specificity: Ubiquitous. Background This gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Submit: Interacts with CNOT7, tissue specificatis factors that have the potential to regulate cell gr	Protein Name	Protein Tob1
FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsTOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1Observed Band40kDCell PathwayCytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmicFunctionfunction:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors., PTM:Phosphorylated on Ser and Thr residues. similarity:Belongs to the BTG familysubunit:Interacts with CNOT7, tissue specificity:Ubiquitous.BackgroundThis gene encodes a member of the transducer of erbB-2 //B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011].	Immunogen	
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic Tissue Specificity Ubiquitous. Function function: Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase. May physically and/or functionally interact with the erbB-2 receptor tyrosine kinase. BTG family., subunit.Interacts with CNOT7., tissue specificity: Ubiquitous. Background This gene encodes a member of the transducer of erbB-2/B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSee, Aug 2011].	Specificity	Phospho-TOB1 (S164) Polyclonal Antibody detects endogenous levels of TOB1 protein only when phosphorylated at S164.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic Tissue Specificity Ubiquitous. Function function:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase receptorsPTM:Phosphorylated on Ser and Thr residues, similarity:Belongs to the BTG family.,subunit:Interacts with CNOT7tissue specificity:Ubiquitous. Background This gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSee, Aug 2011]. matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic Tissue Specificity Ubiquitous. Function function:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase receptors. PTM:Phosphorylated on Ser and Thr residues.,similarity:Belongs to the BTG family.,subunit:Interacts with CNOT7, tissue specificity:Ubiquitous, Background This gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011], matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic Tissue Specificity Ubiquitous. Function function:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase receptors .,PTM:Phosphorylated on Ser and Thr residues, similarity:Belongs to the BTG family.subunit:Interacts with CNOT7.,tissue specificity:Ubiquitous, Background This gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011], matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1 Observed Band 40kD Cell Pathway Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic Tissue Specificity Ubiquitous. Function function:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors., PTM:Phosphorylated on Ser and Thr residues., similarity:Belongs to the BTG family., subunit:Interacts with CNOT7, tissue specificity:Ubiquitous., Background This gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011], matters needing Avoid repeated freezing and thawing!	Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200
Storage Stability-20°C/1 yearSynonymsTOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1Observed Band40kDCell PathwayCytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmicTissue SpecificityUbiquitous.Functionfunction:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors.,PTM:Phosphorylated on Ser and Thr residues., similarity:Belongs to the BTG family.,subunit:Interacts with CNOT7.,tissue specificity:Ubiquitous.,BackgroundThis gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011],matters needingAvoid repeated freezing and thawing!	Concentration	1 mg/ml
SynonymsTOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1Observed Band40kDCell PathwayCytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmicTissue SpecificityUbiquitous.Functionfunction:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors.,PTM:Phosphorylated on Ser and Thr residues.,similarity:Belongs to the BTG family.,subunit:Interacts with CNOT7.,tissue specificity:Ubiquitous.,BackgroundThis gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011],matters needingAvoid repeated freezing and thawing!	Purity	≥90%
Observed Band40kDCell PathwayCytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmicTissue SpecificityUbiquitous.Functionfunction:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors.,PTM:Phosphorylated on Ser and Thr residues.,similarity:Belongs to the BTG family.,subunit:Interacts with CNOT7.,tissue specificity:Ubiquitous.,BackgroundThis gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011],matters needingAvoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell PathwayCytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmicTissue SpecificityUbiquitous.Functionfunction:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors.,PTM:Phosphorylated on Ser and Thr residues.,similarity:Belongs to the BTG family.,subunit:Interacts with CNOT7.,tissue specificity:Ubiquitous.,BackgroundThis gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011],matters needingAvoid repeated freezing and thawing!	Synonyms	TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1
Iate S-phase where more than half of proteins become cytoplasmicTissue SpecificityUbiquitous.Functionfunction:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors.,PTM:Phosphorylated on Ser and Thr residues.,similarity:Belongs to the BTG family.,subunit:Interacts with CNOT7.,tissue specificity:Ubiquitous.,BackgroundThis gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011],matters needingAvoid repeated freezing and thawing!	Observed Band	40kD
Functionfunction:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors.,PTM:Phosphorylated on Ser and Thr residues.,similarity:Belongs to the BTG family.,subunit:Interacts with CNOT7.,tissue specificity:Ubiquitous.,BackgroundThis gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011],matters needingAvoid repeated freezing and thawing!	Cell Pathway	Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic
 kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors.,PTM:Phosphorylated on Ser and Thr residues.,similarity:Belongs to the BTG family.,subunit:Interacts with CNOT7.,tissue specificity:Ubiquitous., Background This gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011], matters needing Avoid repeated freezing and thawing! 	Tissue Specificity	Ubiquitous.
 gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011], matters needing Avoid repeated freezing and thawing! 	Function	kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors.,PTM:Phosphorylated on Ser and Thr residues.,similarity:Belongs to the
	Background	gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by
	matters needing attention	Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com

Ø Website: www.upingBio.com

Usage suggestions

JK

p-TOB1 (S164)

138---100---70---55---40---


35---

15---

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of JK cells using Phospho-TOB1 (S164) Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using TOB1 (Phospho-Ser164) Antibody. The picture on the right is blocked with the phospho peptide.

