

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com




FASTKD1 Polyclonal Antibody


Catalog No	YP-Ab-00396
Isotype	IgG
Reactivity	Human;Mouse;Monkey
Applications	WB;ELISA
Gene Name	FASTKD1
Protein Name	FAST kinase domain-containing protein 1
Immunogen	The antiserum was produced against synthesized peptide derived from human FAKD1. AA range:561-610
Specificity	FASTKD1 Polyclonal Antibody detects endogenous levels of FASTKD1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
. any	290 /0
Storage Stability	-20°C/1 year
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year FASTKD1; KIAA1800; FAST kinase domain-containing protein 1
Storage Stability Synonyms Observed Band	-20°C/1 year FASTKD1; KIAA1800; FAST kinase domain-containing protein 1 97kD Mitochondrion . Preferentially localizes to mitochondrial RNA granules, platforms for post-transcriptional RNA modification and ribosome assembly
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year FASTKD1; KIAA1800; FAST kinase domain-containing protein 1 97kD Mitochondrion . Preferentially localizes to mitochondrial RNA granules, platforms for post-transcriptional RNA modification and ribosome assembly (PubMed:28335001) Expression detected in spleen, thymus, testis, ovary, colon, heart, smooth muscle, kidney, brain, lung, liver and white adipose tissue with highest expression
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year FASTKD1; KIAA1800; FAST kinase domain-containing protein 1 97kD Mitochondrion . Preferentially localizes to mitochondrial RNA granules, platforms for post-transcriptional RNA modification and ribosome assembly (PubMed:28335001) Expression detected in spleen, thymus, testis, ovary, colon, heart, smooth muscle, kidney, brain, lung, liver and white adipose tissue with highest expression in heart. caution:It is uncertain whether Met-1 or Met-15 is the initiator.,similarity:Belongs to
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year FASTKD1; KIAA1800; FAST kinase domain-containing protein 1 97kD Mitochondrion . Preferentially localizes to mitochondrial RNA granules, platforms for post-transcriptional RNA modification and ribosome assembly (PubMed:28335001) Expression detected in spleen, thymus, testis, ovary, colon, heart, smooth muscle, kidney, brain, lung, liver and white adipose tissue with highest expression in heart. caution:It is uncertain whether Met-1 or Met-15 is the initiator.,similarity:Belongs to the FAST kinase family.,similarity:Contains 1 RAP domain., caution:It is uncertain whether Met-1 or Met-15 is the initiator.,similarity:Belongs to



Products Images



Western Blot analysis of various cells using FASTKD1 Polyclonal Antibody




Western blot analysis of lysates from COS7 cells, using FAKD1 Antibody. The lane on the right is blocked with the synthesized peptide.



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com





Western blot analysis of the lysates from HT-29 cells using FAKD1 antibody.