

Website: www.upingBio.com

A-FABP Monoclonal Antibody

Isotype IgG Reactivity Human Applications WB;ELISA Gene Name FABP4 Protein Name Fatty acid-binding protein, adipocyte Immunogen Purified recombinant fragment of A-FABP expressed in E. Coli. Specificity A-FABP Monocional Antibody detects endogenous levels of A-FABP protein. Formulation Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol. Source Monocional, Mouse Purification Affinity purification Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; ALBP; ArBP; Fatty acid-binding protein; ALBP; ArBP; Fatty acid-binding protein to mature of the ligand, a conformation of angle exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export. Cell Pathway Cytoplasm. Nucleus. Depending on the nature of the ligand, a conformation of angle exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export.		
Reactivity Human Applications WB;ELISA Gene Name FABP4 Protein Name Fatty acid-binding protein, adipocyte Immunogen Purified recombinant fragment of A-FABP expressed in E. Coli. Specificity A-FABP Monoclonal Antibody detects endogenous levels of A-FABP protein. Formulation Ascitic fluid containing 0.03% sodium azide, 0.5% BSA, 50%glycerol. Source Monoclonal, Mouse Purification Affinity purification Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein; ALBP; Adipocyte type fatty acid-binding protein is transported into the nucleus. Subject to constitutive nuclear export. Cell Pathway Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export. Function domain;Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior. function: Lipid transport protein in adipocytes. Brids both long chain its insterior. Thaction: Lipid transport protein in stransported into the ealycin	Catalog No	YP-Ab-00595
Applications WB;ELISA Gene Name FABP4 Protein Name Fatty acid-binding protein, adipocyte Immunogen Purified recombinant fragment of A-FABP expressed in E. Coli. Specificity A-FABP Monoclonal Antibody detects endogenous levels of A-FABP protein. Formulation Ascitic fluid containing 0.03% sodium azide, 0.5% BSA, 50%glycerol. Source Monoclonal, Mouse Purification Affinity purification Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4 Observed Band Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior. function: Lipid transport protein in stransported into the calycin superfamily. Fatry acid binding protein in transported into the calyc	Isotype	lgG
Gene Name FABP4 Protein Name Fatty acid-binding protein, adipocyte Immunogen Purified recombinant fragment of A-FABP expressed in E. Coli. Specificity A-FABP Monoclonal Antibody detects endogenous levels of A-FABP protein. Formulation Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol. Source Monoclonal, Mouse Purification Affinity purification Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein; ALBP; Acipocyte-lipid-binding protein 4 Observed Band Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export. Function domin;Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior, function.Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid to their cognate receptors in the nucleussimilarity.Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family, subcellular localization motif and the protein is transported in to	Reactivity	Human
Protein Name Fatty acid-binding protein, adipocyte Protein Name Fatty acid-binding protein, adipocyte Immunogen Purified recombinant fragment of A-FABP expressed in E, Coli. Specificity A-FABP Monoclonal Antibody detects endogenous levels of A-FABP protein. Formulation Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol. Source Monoclonal, Mouse Purification Affinity purification Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4 Observed Band Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export. Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior; function: Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cogna	Applications	WB;ELISA
Immunogen Purified recombinant fragment of A-FABP expressed in E. Coli. Specificity A-FABP Monoclonal Antibody detects endogenous levels of A-FABP protein. Formulation Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol. Source Monoclonal, Mouse Purification Affinity purification Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; Fatty acid-binding protein 4 Observed Band Cytoplasm. Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export. Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior, function.Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long, a conformation chain gexposes a nuclear localization motif and the protein chain chain gexposes a nuclear localization motif and the protein stransported into the calycin superfamily. Fatty-acid binding protein (FABP) family, subcellular location between the to constitutive nucleus. similarity: Belongs to the calycin superfamily. Fatty-acid binding protei	Gene Name	FABP4
Specificity A-FABP Monoclonal Antibody detects endogenous levels of A-FABP protein. Formulation Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol. Source Monoclonal, Mouse Purification Affinity purification Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein. Cell Pathway Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior, function: Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus. subject to constitutive nuclear export. Background FABP4 encodes the fatty acid binding protein formation change exposes a nuclear localization motif and the protein is transported into the nucleus. Sub	Protein Name	Fatty acid-binding protein, adipocyte
Formulation Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol. Source Monoclonal, Mouse Purification Affinity purification Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4 Observed Band Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in fatty acids and retinoic acid to their cognate receptors in the nucleus, similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family, subcellular location their cognate receptors in the nucleus, similarity:Belongs to the calycin superfamily. Fatty-acid binding protein is transported into the nucleus. Subject to constitutive nuclear export. Background FABP4 encodes the fatty acid binding protein in a dipocytes. Fatty acid binding notein is the nucleus. Subject to constitutive nuclear export.	Immunogen	Purified recombinant fragment of A-FABP expressed in E. Coli.
Source Monoclonal, Mouse Purification Affinity purification Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4 Observed Band Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export. Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior. Indicion: Lipid transport protein in adipocytes. Subject to constitutive nucleus. similarity, acids and retinoic acid to their cognate receptors in the nucleus.similarity. Belongs to the calycin superfamily. Fatty-acid binding protein is transported into the nucleus. Subject to constitutive nuclear export. Background FABP4 encodes the fatty acid binding protein found in adipocytes.	Specificity	A-FABP Monoclonal Antibody detects endogenous levels of A-FABP protein.
Purification Affinity purification Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4 Observed Band Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior.,function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity.elongs to the calycin super family. Fatty-acid binding protein (FABP) family. Subcellular localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export. Background FABP4 encodes the fatty acid binding protein in adipocytes. Fatty acid binding protein is the nucleus. Subject to constitutive nuclear export.	Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4 Observed Band Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior., function: Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus., similarity.Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family, subcellular location:Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus, similarity.Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family, subcellular location:Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export, subunit:Homodimer. Interacts with PPARG (By similarity). Monomer.,	Source	Monoclonal, Mouse
applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4 Observed Band Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior, function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family, subcellular location:Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export Background FABP4 encodes the fatty acid binding protein found in adipocytes. Fatty acid binding protein stransported into the nucleus. Subject to constitutive nuclear export, subunit:Homodimer. Interacts with PPARG (By similarity). Monomer.,	Purification	Affinity purification
Purity ≥90% Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4 Observed Band Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export. Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior.,function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.,subcellular localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export., subunit:Homodimer. Interacts with PPARG (By similarity). Monomer., Background FABP4 encodes the fatty acid binding protein found in adipocytes. Fatty acid binding proteins are a family of small. highly conserved, cytoplasmic proteins that	Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Storage Stability -20°C/1 year Synonyms FABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4 Observed Band Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior.,function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family., subcellular location:Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export., subunit:Homodimer. Interacts with PPARG (By similarity). Monomer., Background FABP4 encodes the fatty acid binding protein found in adipocytes. Fatty acid binding proteins are a family of small. highly conserved, cytoplasmic proteins that	Concentration	1 mg/ml
SynonymsFABP4; Fatty acid-binding protein; adipocyte; Adipocyte lipid-binding protein; ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4Observed BandCytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear exportTissue SpecificityUrinary bladder,Functiondomain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior.,function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein is transported into the nucleus. Subject to constitutive nuclear export, subunit:Homodimer. Interacts with PPARG (By similarity). Monomer.,BackgroundFABP4 encodes the fatty acid binding protein found in adipocytes. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that	Purity	≥90%
ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty acid-binding protein 4 Observed Band Cell Pathway Cytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export Tissue Specificity Urinary bladder, Function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior.,function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.,subcellular location:Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export.,subunit:Homodimer. Interacts with PPARG (By similarity). Monomer., Background FABP4 encodes the fatty acid binding protein found in adipocytes. Fatty acid binding protein sare a family of small, highly conserved, cytoplasmic proteins that	Storage Stability	-20°C/1 year
Cell PathwayCytoplasm . Nucleus . Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear exportTissue SpecificityUrinary bladder,Functiondomain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior.,function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.,subcellular 	Synonyms	ALBP; Adipocyte-type fatty acid-binding protein; A-FABP; AFABP; Fatty
 change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export Tissue Specificity Urinary bladder, function domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior.,function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.,subcellular location:Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export.,subunit:Homodimer. Interacts with PPARG (By similarity). Monomer., Background 	Observed Band	
Functiondomain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior.,function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.,subcellular location:Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export.,subunit:Homodimer. Interacts with PPARG (By similarity). Monomer.,BackgroundFABP4 encodes the fatty acid binding protein found in adipocytes. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that	Cell Pathway	change exposes a nuclear localization motif and the protein is transported into the
 its interior.,function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.,subcellular location:Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export.,subunit:Homodimer. Interacts with PPARG (By similarity). Monomer., Background FABP4 encodes the fatty acid binding protein found in adipocytes. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that 	Tissue Specificity	Urinary bladder,
binding proteins are a family of small, highly conserved, cytoplasmic proteins that	Function	its interior.,function:Lipid transport protein in adipocytes. Binds both long chain fatty acids and retinoic acid. Delivers long-chain fatty acids and retinoic acid to their cognate receptors in the nucleus.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.,subcellular location:Depending on the nature of the ligand, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus. Subject to constitutive nuclear export.,subunit:Homodimer. Interacts with PPARG (By
	Background	FABP4 encodes the fatty acid binding protein found in adipocytes. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that bind long-chain fatty acids and other hydrophobic ligands. It is thought that FABPs



UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com

Website: www.upingBio.com roles include fatty acid uptake, transport, and metabolism. [provided by RefSeq,

	Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

