



Dynamin I Monoclonal Antibody

Catalog No	YP-Ab-00610
Isotype	IgG
Reactivity	Human
Applications	WB;IHC;IF;ELISA
Gene Name	DNM1
Protein Name	Dynamin-1
Immunogen	Purified recombinant fragment of human Dynamin I expressed in E. Coli.
Specificity	Dynamin I Monoclonal Antibody detects endogenous levels of Dynamin I protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	WB: 1/500 - 1/2000. IHC: 1/200 - 1/1000. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DNM1; DNM; Dynamin-1
Observed Band	
Cell Pathway	Cytoplasm . Cytoplasm, cytoskeleton . Microtubule-associated.
Tissue Specificity	Brain,Platelet,PNS,
Function	catalytic activity:GTP + H(2)O = GDP + phosphate.,function:Microtubule-associated force-producing protein involved in producing microtubule bundles and able to bind and hydrolyze GTP. Most probably involved in vesicular trafficking processes, in particular endocytosis.,similarity:Belongs to the dynamin family.,similarity:Contains 1 GED domain.,similarity:Contains 1 PH domain.,subcellular location:Microtubule-associated.,subunit:Interacts with CAV1 and SH3GLB1. Binds SH3GL1, SH3GL2 and SH3GL3.,
Background	dynamin 1(DNM1) Homo sapiens This gene encodes a member of the dynamin subfamily of GTP-binding proteins. The encoded protein possesses unique mechanochemical properties used to tubulate and sever membranes, and is involved in clathrin-mediated endocytosis and other vesicular trafficking processes. Actin and other cytoskeletal proteins act as binding partners for the encoded protein, which can also self-assemble leading to stimulation of GTPase activity. More than sixty highly conserved copies of the 3' region of this gene are found elsewhere in the genome, particularly on chromosomes Y and 15.



Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis using Dynamin I Monoclonal Antibody against C6 (1), NIH/3T3 (2), SKN-SH (3), LN18 (4), SHSY5Y (5) cell lysate and rat brain tissues lysate (6).



Immunohistochemistry analysis of paraffin-embedded human lymph tissue (A), glioma tissue (B) and cerebellum tissue (C), showing membrane localization with DAB staining using Dynamin I Monoclonal Antibody.

