



ABCC12 Polyclonal Antibody

Catalog No	YP-Ab-00653
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	ABCC12
Protein Name	Multidrug resistance-associated protein 9
Immunogen	The antiserum was produced against synthesized peptide derived from human MRP9. AA range:691-740
Specificity	ABCC12 Polyclonal Antibody detects endogenous levels of ABCC12 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ABCC12; MRP9; Multidrug resistance-associated protein 9; ATP-binding cassette sub-family C member 12
Observed Band	170kD
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein .
Tissue Specificity	Expressed in testis (at protein level). Widely expressed at low level (PubMed:11483364, PubMed:11688999, PubMed:12011458, PubMed:17472575). Isoform 5 is specifically expressed in brain, testis and breast cancer cells (PubMed:11483364, PubMed:11688999, PubMed:12011458).
Function	developmental stage:Expressed in fetal tissues.,function:Probable transporter.,similarity:Belongs to the ABC transporter family. Conjugate transporter (TC 3.A.1.208) subfamily.,similarity:Contains 2 ABC transmembrane type-1 domains.,similarity:Contains 2 ABC transporter domains.,tissue specificity:Expressed in testis (at protein level). Widely expressed at low level. Isoform 5 is specifically expressed in brain, testis and breast cancer cells.,
Background	This gene is a member of the superfamily of ATP-binding cassette (ABC) transporters and the encoded protein contains two ATP-binding domains and 12 transmembrane regions. ABC proteins transport various molecules across extra- and intracellular membranes. ABC genes are divided into seven distinct subfamilies: ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, and White. This gene



is a member of the MRP subfamily which is involved in multi-drug resistance. This gene and another subfamily member are arranged head-to-tail on chromosome 16q12.1. Increased expression of this gene is associated with breast cancer. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using ABCC12 Polyclonal Antibody diluted at 1:500



Western blot analysis of lysates from 293, HepG2, and HUVEC cells, using MRP9 Antibody. The lane on the right is blocked with the synthesized peptide.