

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



GS28 Polyclonal Antibody

| Catalog No | YP-Ab-00703 |
|--------------------|--|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB;ELISA |
| Gene Name | GOSR1 |
| Protein Name | Golgi SNAP receptor complex member 1 |
| Immunogen | The antiserum was produced against synthesized peptide derived from human GOSR1. AA range:11-60 |
| Specificity | GS28 Polyclonal Antibody detects endogenous levels of GS28 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | GOSR1; GS28; Golgi SNAP receptor complex member 1; 28 kDa Golgi SNARE protein; 28 kDa cis-Golgi SNARE p28; GOS-28 |
| Observed Band | 28kD |
| Cell Pathway | Golgi apparatus membrane; Single-pass type IV membrane protein. Localizes throughout the Golgi apparatus, with lowest levels in the trans-Golgi network (By similarity). Enriched on vesicular components at the terminal rims of the Golgi. Found in Golgi microtubules at low temperature (15 degrees Celsius). |
| Tissue Specificity | Colon carcinoma,Prostate,Testis,Umbilical cord blood, |
| Function | function:Involved in transport from the ER to the Golgi apparatus as well as in intra-Golgi transport.,similarity:Belongs to the GOSR1 family.,subcellular location:Enriched on vesicular components at the terminal rims of the Golgi.,subunit:Interacts with GABARAPL2. Identified in a unique SNARE complex composed of the Golgi SNAREs GOSR2, STX5 and YKT6., |
| Background | This gene encodes a trafficking membrane protein which transports proteins among the endoplasmic reticulum and the Golgi and between Golgi compartments. This protein is considered an essential component of the Golgi SNAP receptor (SNARE) complex. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008], |



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:Upingbio.163.com




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from COLO and K562 cells, using GOSR1 Antibody. The lane on the right is blocked with the synthesized peptide.