



# JMJD1A Monoclonal Antibody

<b>Catalog No</b>	YP-Ab-00995
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human
<b>Applications</b>	WB;IF;ELISA
<b>Gene Name</b>	KDM3A
<b>Protein Name</b>	Lysine-specific demethylase 3A
<b>Immunogen</b>	Purified recombinant fragment of human JMJD1A expressed in E. Coli.
<b>Specificity</b>	JMJD1A Monoclonal Antibody detects endogenous levels of JMJD1A protein.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
<b>Source</b>	Monoclonal, Mouse
<b>Purification</b>	Affinity purification
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	KDM3A; JHDM2A; JMJD1; JMJD1A; KIAA0742; TSGA; Lysine-specific demethylase 3A; JmjC domain-containing histone demethylation protein 2A; Jumonji domain-containing protein 1A
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm . Nucleus . Nuclear in round spermatids. When spermatids start to elongate, localizes to the cytoplasm where it forms distinct foci which disappear in mature spermatozoa (By similarity) .
<b>Tissue Specificity</b>	Adrenal gland,Brain,Fetal kidney,Salivary gland,Testis,
<b>Function</b>	cofactor: Binds 1 Fe(2+) ion per subunit.,domain:Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs are known to mediate the association with nuclear receptors.,domain:The JmjC domain and the C6-type zinc-finger are required for the demethylation activity.,function:Histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a central role in histone code. Preferentially demethylates mono- and dimethylated H3 'Lys-9' residue, with a preference for dimethylated residue, while it has weak or no activity on trimethylated H3 'Lys-9'. Demethylation of Lys residue generates formaldehyde and succinate. Involved in hormone-dependent transcriptional activation, by participating in recruitment to androgen-receptor target genes, resulting in H3 'Lys-9' demethylation and transcriptional activation. Involved in spermatogenesis by regulating expression of target genes such as PRM1 and TMP1



**Background**

This gene encodes a zinc finger protein that contains a jumonji domain and may play a role in hormone-dependent transcriptional activation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2009],


**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western Blot analysis using JMJD1A Monoclonal Antibody against HeLa (1) and HepG2 (2) cell lysate.



Immunofluorescence analysis of Hela cells using JMJD1A Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.