



# LexA DNA Binding Region Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-01195
<b>Isotype</b>	IgG
<b>Reactivity</b>	0
<b>Applications</b>	WB
<b>Gene Name</b>	
<b>Protein Name</b>	
<b>Immunogen</b>	Recombinant Protein of LexA DNA Binding Region
<b>Specificity</b>	The antibody detects endogenous LexA DNA Binding Region protein.
<b>Formulation</b>	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB: 1:10000-50000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	YM3412
<b>Observed Band</b>	
<b>Cell Pathway</b>	
<b>Tissue Specificity</b>	
<b>Function</b>	
<b>Background</b>	LexA (EC 3. 1. 8) is a DNA binding protein, is associated with SOS response of cyanobacteria as well as other regulatory networks. Has been shown to directly regulate genes involved in carbon assimilation or controlled by carbon availability. In Synechocystis PCC 6803 LexA is not localised to the thylakoid membranes but to the inner region of the cytoplasm.
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images



Western blot analysis of Recombinant LexA protein using LexA DNA Binding Region Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000