



# HMG-17 (phospho Ser29) Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-01407
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	IHC;IF;ELISA
<b>Gene Name</b>	HMGN2
<b>Protein Name</b>	Non-histone chromosomal protein HMG-17
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human HMG17 around the phosphorylation site of Ser29. AA range:1-50
<b>Specificity</b>	Phospho-HMG-17 (S29) Polyclonal Antibody detects endogenous levels of HMG-17 protein only when phosphorylated at S29.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	IHC: 1/100 - 1/300. ELISA: 1/5000.. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	HMGN2; HMG17; Non-histone chromosomal protein HMG-17; High mobility group nucleosome-binding domain-containing protein 2
<b>Observed Band</b>	15-17kD
<b>Cell Pathway</b>	Nucleus . Cytoplasm . Cytoplasmic enrichment upon phosphorylation.
<b>Tissue Specificity</b>	Brain,Colon,Eye,Lung,Lymph,Mammary gland,Muscle,Peripheral
<b>Function</b>	function: Binds to the inner side of the nucleosomal DNA thus altering the interaction between the DNA and the histone octamer. May be involved in the process which maintains transcribable genes in a unique chromatin conformation.,mass spectrometry: PubMed:10739259,PTM:Phosphorylation favors cytoplasmic localization.,similarity:Belongs to the HMGN family.,subcellular location:Cytoplasmic enrichment upon phosphorylation.,
<b>Background</b>	high mobility group nucleosomal binding domain 2(HMGN2) Homo sapiens The protein encoded by this gene binds nucleosomal DNA and is associated with transcriptionally active chromatin. Along with a similar protein, HMGN1, the encoded protein may help maintain an open chromatin configuration around transcribable genes. The protein has also been found to have antimicrobial activity against bacteria, viruses and fungi. [provided by RefSeq, Oct 2014],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.