



# TFIIIA Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-02108
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	GTF3A
<b>Protein Name</b>	Transcription factor IIIA
<b>Immunogen</b>	Synthesized peptide derived from the Internal region of human TFIIIA.
<b>Specificity</b>	TFIIIA Polyclonal Antibody detects endogenous levels of TFIIIA protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	GTF3A; Transcription factor IIIA; TFIIIA
<b>Observed Band</b>	41kD
<b>Cell Pathway</b>	Nucleus.
<b>Tissue Specificity</b>	Ubiquitous.
<b>Function</b>	caution:It is uncertain whether Met-1 is the initiator. Based on the lack of an in-frame AUG codon, mammalian TFIIIA may be translated from this non-AUG initiation site, which has a good Kozak context and which is well conserved among mammals.,function:Interacts with the internal control region (ICR) of approximately 50 bases within the 5S RNA genes, is required for correct transcription of these genes by RNA polymerase III. Also binds the transcribed 5S RNA's. May initiate transcription of the 5S ribosomal RNA gene and maintain the stability of transcription of other genes.,sequence caution:Translation N-terminally shortened.,similarity:Contains 9 C2H2-type zinc fingers.,tissue specificity:Ubiquitous.,
<b>Background</b>	The product of this gene is a zinc finger protein with nine Cis[2]-His[2] zinc finger domains. It functions as an RNA polymerase III transcription factor to induce transcription of the 5S rRNA genes. The protein binds to a 50 bp internal promoter in the 5S genes called the internal control region (ICR), and nucleates formation of a stable preinitiation complex. This complex recruits the TFIIC and TFIIB



transcription factors and RNA polymerase III to form the complete transcription complex. The protein is thought to be translated using a non-AUG translation initiation site in mammals based on sequence analysis, protein homology, and the size of the purified protein. [provided by RefSeq, Jul 2008],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4°over night