



# PKM2 mouse mAb

<b>Catalog No</b>	YP-Ab-02362
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat;Monkey
<b>Applications</b>	WB;IF
<b>Gene Name</b>	pkm2
<b>Protein Name</b>	
<b>Immunogen</b>	Purified recombinant human PKM2 protein fragments expressed in E.coli.
<b>Specificity</b>	This antibody detects endogenous levels of PKM2 and does not cross-react with related proteins.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	wb 1:1000 icc 1:400. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	CTHBP;Cytosolic thyroid hormone binding protein;Cytosolic thyroid hormone-binding protein;KPYM_HUMAN;MGC3932;OIP 3;OIP-3;OIP3;OPA interacting protein 3;Opa-interacting protein 3;p58;PK muscle type;PK, muscle type;PK2;PK3;PKM;PKM2;pykm;Pyruvate kinase 2/3;Pyruvate kinase 3;Pyruvate kinase isozymes M1/M2;Pyruvate kinase muscle;Pyruvate kinase muscle isozyme;pyruvate kinase PKM;Pyruvate kinase, muscle 2;TCB;THBP1;Thyroid hormone binding protein 1;Thyroid hormone binding protein cytosolic;Thyroid hormone-binding protein 1;Tumor M2 PK;Tumor M2-PK.
<b>Observed Band</b>	60kD
<b>Cell Pathway</b>	[Isoform M2]: Cytoplasm . Nucleus . Translocates to the nucleus in response to various signals, such as EGF receptor activation or apoptotic stimuli (PubMed:17308100, PubMed:22056988, PubMed:24120661). Nuclear translocation is promoted by acetylation by EP300 (PubMed:24120661). Deacetylation by SIRT6 promotes its nuclear export in a process dependent of XPO4, thereby suppressing its ability to activate transcription and promote tumorigenesis (PubMed:26787900). .; [Isoform M1]: Cytoplasm .
<b>Tissue Specificity</b>	[Isoform M2]: Specifically expressed in proliferating cells, such as embryonic stem cells, embryonic carcinoma cells, as well as cancer cells. ; [Isoform M1]: Expressed in adult tissues (PubMed:18337823). Not expressed in tumor cells




(PubMed:18337823).


<b>Function</b>	<p>catalytic activity:ATP + pyruvate = ADP + phosphoenolpyruvate.,cofactor:Divalent metal cations.,cofactor:Magnesium.,cofactor:Potassium.,enzyme regulation:Isoform M2 is allosterically activated by D-fructose 1,6-biphosphate (FBP). Inhibited by oxalate and 3,3',5-triiodo-L-thyronine (T3).,function:Glycolytic enzyme that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate (PEP) to ADP, generating ATP.,miscellaneous:There are 4 isozymes of pyruvate kinase in mammals: L, R, M1 and M2. L type is major isozyme in the liver, R is found in red cells, M1 is the main form in muscle, heart and brain, and M2 is found in early fetal tissues as well as in most cancer cells.,online information:Pyruvate kinase entry,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 5/5.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Be</p>
<b>Background</b>	<p>This gene encodes a protein involved in glycolysis. The encoded protein is a pyruvate kinase that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP, generating ATP and pyruvate. This protein has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. This protein has been found to bind Opa protein, a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells, suggesting a role of this protein in bacterial pathogenesis. Several alternatively spliced transcript variants encoding a few distinct isoforms have been reported. [provided by RefSeq, May 2011],</p>
<b>matters needing attention</b>	<p>Avoid repeated freezing and thawing!</p>
<b>Usage suggestions</b>	<p>This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.</p>



## Products Images



Western blot detection of PKM2 in COS7,PC12,C6,3T3 and HeLa cell lysates using PKM2 mouse mAb (1:1000 diluted).Predicted band size:60KDa.Observed band size:60KDa.



Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using anti-PKM2 mouse mAb (dilution 1:400).