



PDK1 (Phospho Thr338) rabbit pAb

Catalog No	YP-Ab-02442
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB; ELISA;IHC
Gene Name	PDK1 PDHK1
Protein Name	PDHK1 (Phospho Thr338)
Immunogen	Synthesized peptide derived from human PDHK1 (Phospho Thr338)
Specificity	This antibody detects endogenous levels of Human,Mouse,Rat PDHK1 (Phospho Thr338)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	[Pyruvate dehydrogenase [lipoamide]] kinase isozyme 1, mitochondrial (EC 2.7.11.2;Pyruvate dehydrogenase kinase isoform 1)
Observed Band	50kD
Cell Pathway	Mitochondrion matrix .
Tissue Specificity	Expressed predominantly in the heart. Detected at lower levels in liver, skeletal muscle and pancreas.
Function	catalytic activity:ATP + [pyruvate dehydrogenase (acetyl-transferring)] = ADP + [pyruvate dehydrogenase (acetyl-transferring)] phosphate.,function:Inhibits the mitochondrial pyruvate dehydrogenase complex by phosphorylation of the E1 alpha subunit, thus contributing to the regulation of glucose metabolism.,similarity:Belongs to the PDK/BCKDK protein kinase family.,similarity:Contains 1 histidine kinase domain.,tissue specificity:Expressed predominantly in the heart.,
Background	Pyruvate dehydrogenase (PDH) is a mitochondrial multienzyme complex that catalyzes the oxidative decarboxylation of pyruvate and is one of the major enzymes responsible for the regulation of homeostasis of carbohydrate fuels in mammals. The enzymatic activity is regulated by a phosphorylation/dephosphorylation cycle. Phosphorylation of PDH by a specific pyruvate dehydrogenase kinase (PDK) results in inactivation. Multiple alternatively spliced transcript variants have been found for this gene. [provided



by RefSeq, Jun 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).