

Tel: 400-999-8863
■ Emall:Upingbio.163.com



FX Polyclonal Antibody

| Catalog No | YP-Ab-02637 |
|--------------------|--|
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB;ELISA |
| Gene Name | TSTA3 |
| Protein Name | GDP-L-fucose synthase |
| Immunogen | The antiserum was produced against synthesized peptide derived from human TSTA3. AA range:221-270 |
| Specificity | FX Polyclonal Antibody detects endogenous levels of FX protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | TSTA3; SDR4E1; GDP-L-fucose synthase; GDP-4-keto-6-deoxy-D-mannose-3; 5-epimerase-4-reductase; Protein FX; Red cell NADP(H)-binding protein; Short-chain dehydrogenase/reductase family 4E member 1 |
| Observed Band | 40kD |
| Cell Pathway | cytoplasm,cytosol,extracellular exosome, |
| Tissue Specificity | Erythrocyte,Lung,Placenta,Skin, |
| Function | catalytic activity:GDP-L-fucose + NADP(+) = GDP-4-dehydro-6-deoxy-D-mannose + NADPH.,function:Two step NADP-dependent conversion of GDP-4-dehydro-6-deoxy-D-mannose to GDP-fucose, involving an epimerase and a reductase reaction.,pathway:Nucleotide-sugar biosynthesis; GDP-L-fucose biosynthesis via de novo pathway; GDP-L-fucose from GDP-D-mannose: step 2/2.,similarity:Belongs to the fucose synthetase family.,subunit:Homodimer., |
| Background | Tissue specific transplantation antigen P35B is a NADP(H)-binding protein. It catalyze the two-step epimerase and the reductase reactions in GDP-D-mannose metabolism, converting GDP-4-keto-6-D-deoxymannose to GDP-L-fucose. GDP-L-fucose is the substrate of several fucosyltransferases involved in the expression of many glycoconjugates, including blood group ABH antigens and |



UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 💌 Emall:Upingbio.163.com



developmental adhesion antigens. Mutations in this gene may cause leukocyte adhesion deficiency, type II. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from rat kidney and rat muscle cells, using TSTA3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using TSTA3 antibody.