



# GSTT1/4 Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-02649
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	GSTT1/GSTT4
<b>Protein Name</b>	Glutathione S-transferase theta-1/Glutathione S-transferase theta-4
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human GSTT1/4. AA range:10-59
<b>Specificity</b>	GSTT1/4 Polyclonal Antibody detects endogenous levels of GSTT1/4 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	GSTT1; Glutathione S-transferase theta-1; GST class-theta-1; Glutathione transferase T1-1; GSTT4; Glutathione S-transferase theta-4; GST class-theta-4
<b>Observed Band</b>	30kD
<b>Cell Pathway</b>	Cytoplasm.
<b>Tissue Specificity</b>	Found in erythrocyte. Expressed at low levels in liver. In lung, expressed at low levels in Clara cells and ciliated cells at the alveolar/bronchiolar junction. Absent from epithelial cells of larger bronchioles.
<b>Function</b>	catalytic activity:RX + glutathione = HX + R-S-glutathione.,function:Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. Acts on 1,2-epoxy-3-(4-nitrophenoxy)propane, phenethylisothiocyanate 4-nitrobenzyl chloride and 4-nitrophenethyl bromide. Displays glutathione peroxidase activity with cumene hydroperoxide.,online information:The Singapore human mutation and polymorphism database,polymorphism:The GSTT1 gene is absent from 38% of the population. The presence or absence of the GSTT1 gene is coincident with the conjugator (GSST1+) and non-conjugator (GSTT1-) phenotypes respectively. The GSTT1+ phenotype can catalyze the glutathione conjugation of dichloromethane.,similarity:Belongs to the GST superfamily. Theta family.,similarity:Contains 1 GST C-terminal domain.,similarity:Contains 1 GST N-terminal domain.,subunit:Homodimer.,tissue s



### Background

The protein encoded by this gene, glutathione S-transferase (GST) theta 1 (GSTT1), is a member of a superfamily of proteins that catalyze the conjugation of reduced glutathione to a variety of electrophilic and hydrophobic compounds. Human GSTs can be divided into five main classes: alpha, mu, pi, theta, and zeta. The theta class includes GSTT1, GSTT2, and GSTT2B. GSTT1 and GSTT2/GSTT2B share 55% amino acid sequence identity and may play a role in human carcinogenesis. The GSTT1 gene is haplotype-specific and is absent from 38% of the population. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Sep 2015],


### matters needing attention

Avoid repeated freezing and thawing!


### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using GSTT1/4 Polyclonal Antibody



Western blot analysis of lysates from Jurkat cells, using GSTT1/4 Antibody. The lane on the right is blocked with the synthesized peptide.