

Website: www.upingBio.com

## **USP53 Polyclonal Antibody**

Catalog No	YP-Ab-02843
Isotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	USP53
Protein Name	Inactive ubiquitin carboxyl-terminal hydrolase 53
Immunogen	The antiserum was produced against synthesized peptide derived from human USP53. AA range:951-1000
Specificity	USP53 Polyclonal Antibody detects endogenous levels of USP53 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Purification Dilution	
	affinity-chromatography using epitope-specific immunogen.
Dilution	affinity-chromatography using epitope-specific immunogen. WB 1:500-2000 IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200
Dilution Concentration	affinity-chromatography using epitope-specific immunogen. WB 1:500-2000 IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200 1 mg/ml
Dilution Concentration Purity	affinity-chromatography using epitope-specific immunogen. WB 1:500-2000 IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200 1 mg/ml ≥90%
Dilution Concentration Purity Storage Stability	affinity-chromatography using epitope-specific immunogen. WB 1:500-2000 IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200 1 mg/ml ≥90% -20°C/1 year USP53; KIAA1350; Inactive ubiquitin carboxyl-terminal hydrolase 53; Inactive
Dilution Concentration Purity Storage Stability Synonyms	affinity-chromatography using epitope-specific immunogen. WB 1:500-2000 IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200 1 mg/ml ≥90% -20°C/1 year USP53; KIAA1350; Inactive ubiquitin carboxyl-terminal hydrolase 53; Inactive
Dilution Concentration Purity Storage Stability Synonyms Observed Band	affinity-chromatography using epitope-specific immunogen. WB 1:500-2000 IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200 1 mg/ml ≥90% -20°C/1 year USP53; KIAA1350; Inactive ubiquitin carboxyl-terminal hydrolase 53; Inactive ubiquitin-specific peptidase 53
Dilution Concentration Purity Storage Stability Synonyms Observed Band Cell Pathway	affinity-chromatography using epitope-specific immunogen. WB 1:500-2000 IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200 1 mg/ml ≥90% -20°C/1 year USP53; KIAA1350; Inactive ubiquitin carboxyl-terminal hydrolase 53; Inactive ubiquitin-specific peptidase 53 Cell junction, tight junction .
Dilution Concentration Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	affinity-chromatography using epitope-specific immunogen. WB 1:500-2000 IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200 1 mg/ml ≥90% -20°C/1 year USP53; KIAA1350; Inactive ubiquitin carboxyl-terminal hydrolase 53; Inactive ubiquitin-specific peptidase 53 Cell junction, tight junction . Expressed predominantly in skeletal muscle and heart. caution:Although the active site residues are conserved, lacks the conserved His residue which is normally found 9 residues before the catalytic His.,function:Has no peptidase activity.,similarity:Belongs to the peptidase C19 family.,tissue



## UpingBio technology Co.,Ltd


🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com

Website: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using USP53 Antibody. The picture on the right is blocked with the synthesized peptide.