



Pinch-1 Monoclonal Antibody

Catalog No	YP-Ab-03414
Isotype	IgG
Reactivity	Human
Applications	WB;IF;FCM;ELISA
Gene Name	LIMS1
Protein Name	LIM and senescent cell antigen-like-containing domain protein 1
Immunogen	Purified recombinant fragment of human Pinch-1 expressed in E. Coli.
Specificity	Pinch-1 Monoclonal Antibody detects endogenous levels of Pinch-1 protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. Flow cytometry: 1/200 - 1/400. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	LIMS1; PINCH; PINCH1; LIM and senescent cell antigen-like-containing domain protein 1; Particularly interesting new Cys-His protein 1; PINCH-1; Renal carcinoma antigen NY-REN-48
Observed Band	
Cell Pathway	Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side.
Tissue Specificity	Expressed in most tissues except in the brain.
Function	function:Effector of integrin and growth factor signaling, coupling surface receptors to downstream signaling molecules involved in the regulation of cell survival, cell proliferation and cell differentiation. Focal adhesion protein part of the complex ILK-PINCH. This complex is considered to be one of the convergence points of integrin- and growth factor-signaling pathway.,similarity:Contains 5 LIM zinc-binding domains.,subunit:Interacts (via LIM zinc-binding 5) with TGFB111 (By similarity). Interacts (via LIM zinc-binding 1) with ILK. Interacts with SH3/SH2 adapter NCK2.,tissue specificity:In most tissues, except in the brain.,
Background	The protein encoded by this gene is an adaptor protein which contains five LIM domains, or double zinc fingers. The protein is likely involved in integrin signaling through its LIM domain-mediated interaction with integrin-linked kinase, found in focal adhesion plaques. It is also thought to act as a bridge linking integrin-linked



kinase to NCK adaptor protein 2, which is involved in growth factor receptor kinase signaling pathways. Its localization to the periphery of spreading cells also suggests that this protein may play a role in integrin-mediated cell adhesion or spreading. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2010],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis using Pinch-1 Monoclonal Antibody against A549 (1), Jurkat (2), and HeLa (3) cell lysate.



Immunofluorescence analysis of HepG2 cells using Pinch-1 Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Flow cytometric analysis of HeLa cells using Pinch-1 Monoclonal Antibody (blue) and negative control (red).

