

**(** Tel: 400-999-8863 ■ Emall:Upingbio.163.com



# Hsp70(C-term) mouse mAb

Catalog No	YP-Ab-03451
Isotype	IgG
Reactivity	Human
Applications	WB;IP
Gene Name	hspa1a
Protein Name	
Immunogen	Purified recombinant human Hsp70(C-terminus) protein fragments expressed in E.coli.
Specificity	This antibody detects endogenous levels of Hsp70(C-terminus) and does not cross-react with related proteins.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	wb 1:1000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DAQB 147D11.1 001;FLJ54303;FLJ54370;FLJ54392;FLJ54408;FLJ75127;Heat shock 70 kDa protein 1;Heat shock 70 kDa protein 1/2;Heat shock 70 kDa protein 1A/1B;heat shock 70kDa protein 1A;Heat shock 70kDa protein 1B;Heat shock induced protein;heat shock protein 70;HSP70 1;HSP70 2;HSP70-1/HSP70-2;HSP70-1A;HSP70.1;HSP70.1/HSP70.2;HSP70I;HSP71_H UMAN;HSP72; HSPA1;HSPA1A;HSPA1B;XXbac BCX40G17.3 001.
Observed Band	70kD
Cell Pathway	Cytoplasm . Nucleus . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Secreted . Localized in cytoplasmic mRNP granules containing untranslated mRNAs.
Tissue Specificity	Brain,Cajal-Retzius cell,Embryonic kidney,Epithelium,Fetal
Function	function:In cooperation with other chaperones, Hsp70s stabilize preexistent proteins against aggregation and mediate the folding of newly translated polypeptides in the cytosol as well as within organelles. These chaperones participate in all these processes through their ability to recognize nonnative conformations of other proteins. They bind extended peptide segments with a net hydrophobic character exposed by polypeptides during translation and membrane translocation, or following stress-induced damage. In case of rotavirus A infection,



### UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 💌 Emall:Upingbio.163.com



serves as a post-attachment receptor for the virus to facilitate entry into the cell.,induction:By heat shock.,similarity:Belongs to the heat shock protein 70 family.,subunit:HSPA1B is found in a sperm-specific complex with CATSPER1 and CATSPERB (By similarity). Interacts with TSC2. Interacts with IRAK1BP1.,tissue specificity:HSPA1B is testis-specific

#### **Background**

This intronless gene encodes a 70kDa heat shock protein which is a member of the heat shock protein 70 family. In conjuction with other heat shock proteins, this protein stabilizes existing proteins against aggregation and mediates the folding of newly translated proteins in the cytosol and in organelles. It is also involved in the ubiquitin-proteasome pathway through interaction with the AU-rich element RNA-binding protein 1. The gene is located in the major histocompatibility complex class III region, in a cluster with two closely related genes which encode similar proteins. [provided by RefSeq, Jul 2008],


## matters needing attention

Avoid repeated freezing and thawing!


### **Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### **Products Images**



Western blot detection of Hsp70(C-terminus) in Hela and HsHela cell lysates using Hsp70(C-terminus) mouse mAb (1:1000 diluted).Predicted band size: 70KDa.Observed band size: 70KDa.



Immunoprecipitation analysis of Hela cell lysates using Hsp70 (C-terminus) mouse mAb.