

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Dvl-3 Polyclonal Antibody

Catalog No YP-Ab-03830 Isotype IgG Reactivity Human;Mouse Applications WB;IHC;IF;ELISA Gene Name DVL3 Protein Name Segment polarity protein dishevelled homolog DVL-3 Immunogen The antiserum was produced against synthesized peptide derived from human DVL3. AA range:326-375 Specificity DVI-3 Polyclonal Antibody detects endogenous levels of DVI-3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm Cell Pathway Cytoplasm . Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, Uniteriactive (via the PDZ domain) with the C-terminal regi		
Reactivity Human:Mouse Applications WB:IHC;IF;ELISA Gene Name DVL3 Protein Name Segment polarity protein dishevelled homolog DVL-3 Immunogen The antiserum was produced against synthesized peptide derived from human DVL3. AA range:326-375 Specificity DvI-3 Polyclonal Artibody detects endogenous levels of DvI-3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20*C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm Cell Pathway Cytoplasm Tissue Specificity Bone marrow, Brain, Hippocampus, Placenta, Uterus, function. May play a role in the signal transduction pathway mediated by multiple Wnt genes, similarity: Contains 1 DIX domain, similarity: Contains 1 DEP domains ywith the C-termi	Catalog No	YP-Ab-03830
Applications WB;IHC;IF;ELISA Gene Name DVL3 Protein Name Segment polarity protein dishevelled homolog DVL-3 Immunogen The antiserum was produced against synthesized peptide derived from human DVL3. AA range:326-375 Specificity Dvl-3 Polyclonal Antibody detects endogenous levels of Dvl-3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm. Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, Function function:May play a role in the signal transduction pathway mediated by multiple Wnt genes, similarity,Contains 1 DEP domain, similarity.Contains 1 DEP domain, similarity.Contains 1 DEP (DHR) domain, similarity.Contains 1 DEP (DHR) domain, similarity.Contains 1 DIX domain, similarity.Contains 1 DX domain, similarity.Contains 1 DX domain, similarity.Contains 1 DX domain, similarity.Contains 1 DX domain may inhibit this interaction. Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that requilates cell proliferation. [provided by	Isotype	IgG
Gene Name DVL3 Protein Name Segment polarity protein dishevelled homolog DVL-3 Immunogen The antiserum was produced against synthesized peptide derived from human DVL3. AA range:326-375 Specificity DVI-3 Polyclonal Antibody detects endogenous levels of DVI-3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm. Tissue Specificity Bone marrow, Brain, Hippocampus, Placenta, Uterus, function: May play a role in the signal transduction pathway mediated by multiple Wnt genes, similarity: Belongs to the DSH family, similarity: Contains 1 DEP domain, similarity Contains 1 DEP domain, similarity Contains 1 DEP domain, subunit-interacts (via the PDZ domain) with the C-terminal regions of VANGL 1 and VANGL 2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DI	Reactivity	Human;Mouse
Protein Name Segment polarity protein dishevelled homolog DVL-3	Applications	WB;IHC;IF;ELISA
Immunogen The antiserum was produced against synthesized peptide derived from human DVL3. AA range:326-375 Specificity DVI-3 Polyclonal Antibody detects endogenous levels of DVI-3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm . Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, Function function:May play a role in the signal transduction pathway mediated by multiple Wnt genes, similarity.Belongs to the DSH family, similarity.Contains 1 DEP domain, similarity.Contains 1 DEP domain, similarity.Contains 1 DIX domain, similarity.Contains 1 PDZ (DHR) domain, similarity.Contains 1 DIX domain, similarity.Contains 1 PDZ (DHR) domain, subunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL 1 and VANGL 2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may in	Gene Name	DVL3
DVL3. AA range:326-375 Specificity DVI-3 Polyclonal Antibody detects endogenous levels of DVI-3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, Function function:May play a role in the signal transduction pathway mediated by multiple Wnt genes, similarity:Belongs to the DSH family, similarity:Contains 1 DEP domain, similarity:Contains 1 DDZ (DHR) domain, similarity:Contains 1 DIX domain, similarity:Contains 1 PDZ (DHR) domain, similarity:Contains 1 DPD (domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the PDZ domain) with the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophilia dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Protein Name	Segment polarity protein dishevelled homolog DVL-3
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm . Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, function:May play a role in the signal transduction pathway mediated by multiple Wnt genes. similarity:Belongs to the DSH family. similarity:Contains 1 DEP domain. subunit:Interacts (via the PD2 domain) internal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PD2 and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Immunogen	
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm . Tissue Specificity Bone marrow, Brain, Hippocampus, Placenta, Uterus, Function function: May play a role in the signal transduction pathway mediated by multiple Wnt genes., similarity: Contains 1 DEP domain, similarity: Contains 1 DEP domain, similarity: Contains 1 DEP domain, similarity: Contains 1 DEV domain, similarity:	Specificity	Dvl-3 Polyclonal Antibody detects endogenous levels of Dvl-3 protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm. Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, function:May play a role in the signal transduction pathway mediated by multiple Wnt genes.,similarity:Belongs to the DSH family.,similarity:Contains 1 DEP domain.,similarity:Contains 1 DEP domain.,similarity:Contains 1 DEP domain.,similarity:Contains 1 PDZ (DHR) domain.,subunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm . Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, Function function:May play a role in the signal transduction pathway mediated by multiple Wnt genes, similarity:Contains 1 DEP domainsimilarity:Contains 1 DEP domainsubunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm . Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, Function function:May play a role in the signal transduction pathway mediated by multiple Wnt genes, similarity:Belongs to the DSH family, similarity:Contains 1 DEP domain, similarity:Contains 1 DIX domain, similarity:Contains 1 PDZ (DHR) domain, subunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction. Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm . Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, Function function:May play a role in the signal transduction pathway mediated by multiple Wnt genes.,similarity:Belongs to the DSH family.,similarity:Contains 1 DEP domain.,similarity:Contains 1 DIX domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 poly (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200
Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm . Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, function:May play a role in the signal transduction pathway mediated by multiple Wnt genes.,similarity:Belongs to the DSH family.,similarity:Contains 1 DEP domain.,similarity:Contains 1 DIX domain, similarity:Contains 1 DDZ (DHR) domain.,subunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Concentration	1 mg/ml
Synonyms DVL3; KIAA0208; Segment polarity protein dishevelled homolog DVL-3; Dishevelled-3; DSH homolog 3 Observed Band Cell Pathway Cytoplasm . Bone marrow,Brain,Hippocampus,Placenta,Uterus, function:May play a role in the signal transduction pathway mediated by multiple Wnt genes.,similarity:Belongs to the DSH family.,similarity:Contains 1 DEP domain.,similarity:Contains 1 DIX domain.,similarity:Contains 1 PDZ (DHR) domain.,subunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Purity	≥90%
Observed Band Cell Pathway Cytoplasm . Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, function:May play a role in the signal transduction pathway mediated by multiple Wnt genes.,similarity:Belongs to the DSH family.,similarity:Contains 1 DEP domain.,similarity:Contains 1 DIX domain.,similarity:Contains 1 PDZ (DHR) domain.,subunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Bone marrow,Brain,Hippocampus,Placenta,Uterus, function function:May play a role in the signal transduction pathway mediated by multiple Wnt genes.,similarity:Belongs to the DSH family.,similarity:Contains 1 DEP domain.,similarity:Contains 1 DIX domain.,similarity:Contains 1 PDZ (DHR) domain.,subunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Synonyms	
Tissue Specificity Bone marrow,Brain,Hippocampus,Placenta,Uterus, function:May play a role in the signal transduction pathway mediated by multiple Wnt genes.,similarity:Belongs to the DSH family.,similarity:Contains 1 DEP domain.,similarity:Contains 1 DIX domain.,similarity:Contains 1 PDZ (DHR) domain.,subunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Observed Band	
Function function:May play a role in the signal transduction pathway mediated by multiple Wnt genes.,similarity:Belongs to the DSH family.,similarity:Contains 1 DEP domain.,similarity:Contains 1 DIX domain.,similarity:Contains 1 PDZ (DHR) domain.,subunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Cell Pathway	Cytoplasm .
Wnt genes., similarity: Belongs to the DSH family., similarity: Contains 1 DEP domain., similarity: Contains 1 DIX domain., similarity: Contains 1 PDZ (DHR) domain., subunit: Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this interaction., Background This gene is a member of a multi-gene family which shares strong similarity with the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Tissue Specificity	Bone marrow,Brain,Hippocampus,Placenta,Uterus,
the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by	Function	Wnt genes., similarity:Belongs to the DSH family., similarity:Contains 1 DEP domain., similarity:Contains 1 DIX domain., similarity:Contains 1 PDZ (DHR) domain., subunit:Interacts (via the PDZ domain) with the C-terminal regions of VANGL1 and VANGL2 (By similarity). Interacts (via the region containing both the PDZ and DEP domains) with LRRFIP2; the DIX domain may inhibit this
	Background	the Drosophila dishevelled gene, dsh. The Drosophila dishevelled gene encodes a cytoplasmic phosphoprotein that regulates cell proliferation. [provided by



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🗷 Emall:Upingbio.163.com




matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images



Western blot analysis of DVL3 Antibody. The lane on the right is blocked with the DVL3 peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using DVL3 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from Jurkat cells using DVL3 antibody.