



# Factor XIII B Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-03866
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	F13B
<b>Protein Name</b>	Coagulation factor XIII B chain
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human F13B. AA range:61-110
<b>Specificity</b>	Factor XIII B Polyclonal Antibody detects endogenous levels of Factor XIII B protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	F13B; Coagulation factor XIII B chain; Fibrin-stabilizing factor B subunit; Protein-glutamine gamma-glutamyltransferase B chain; Transglutaminase B chain
<b>Observed Band</b>	80kD
<b>Cell Pathway</b>	Secreted .
<b>Tissue Specificity</b>	Heart,Liver,Plasma,
<b>Function</b>	disease:Defects in F13B can result in a lifelong bleeding tendency, defective wound healing, and habitual abortion.,function:The B chain of factor XIII is not catalytically active, but is thought to stabilize the A subunits and regulate the rate of transglutaminase formation by thrombin.,online information:Factor XIII entry,online information:The Singapore human mutation and polymorphism database,similarity:Contains 10 Sushi (CCP/SCR) domains.,subunit:Tetramer of two A chains and two B chains.,
<b>Background</b>	This gene encodes coagulation factor XIII B subunit. Coagulation factor XIII is the last zymogen to become activated in the blood coagulation cascade. Plasma factor XIII is a heterotetramer composed of 2 A subunits and 2 B subunits. The A subunits have catalytic function, and the B subunits do not have enzymatic activity and may serve as a plasma carrier molecules. Platelet factor XIII is comprised



only of 2 A subunits, which are identical to those of plasma origin. Upon activation by the cleavage of the activation peptide by thrombin and in the presence of calcium ion, the plasma factor XIII dissociates its B subunits and yields the same active enzyme, factor XIIIa, as platelet factor XIII. This enzyme acts as a transglutaminase to catalyze the formation of gamma-glutamyl-epsilon-lysine crosslinking between fibrin molecules, thus stabilizing the fibrin clot. Factor XIII deficiency is classi


**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


**Products Images**



Western Blot analysis of COLO cells using Factor XIII B Polyclonal Antibody diluted at 1:1000



Immunofluorescence analysis of HUVEC cells, using F13B Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human heart tissue, using F13B Antibody. The picture on the right is blocked with the synthesized peptide.