



# FT $\beta$ Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-03879
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	FNTB
<b>Protein Name</b>	Protein farnesyltransferase subunit beta
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human FNTB. AA range:255-304
<b>Specificity</b>	FT $\beta$ Polyclonal Antibody detects endogenous levels of FT $\beta$ protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	$\geq 90\%$
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	FNTB; Protein farnesyltransferase subunit beta; FTase-beta; CAAX farnesyltransferase subunit beta; Ras proteins prenyltransferase subunit beta
<b>Observed Band</b>	49kD
<b>Cell Pathway</b>	cytosol,microtubule associated complex,protein farnesyltransferase complex,
<b>Tissue Specificity</b>	Neuroblastoma,Placenta,Retina,
<b>Function</b>	catalytic activity:Farnesyl diphosphate + protein-cysteine = S-farnesyl protein + diphosphate.,cofactor:Binds 1 zinc ion per subunit.,function:Catalyzes the transfer of a farnesyl moiety from farnesyl pyrophosphate to a cysteine at the fourth position from the C-terminus of several proteins. The beta subunit is responsible for peptide-binding.,similarity:Belongs to the protein prenyltransferase subunit beta family.,similarity:Contains 5 PFTB repeats.,subunit:Heterodimer of an alpha and a beta subunit.,
<b>Background</b>	catalytic activity:Farnesyl diphosphate + protein-cysteine = S-farnesyl protein + diphosphate.,cofactor:Binds 1 zinc ion per subunit.,function:Catalyzes the transfer of a farnesyl moiety from farnesyl pyrophosphate to a cysteine at the fourth position from the C-terminus of several proteins. The beta subunit is responsible for peptide-binding.,similarity:Belongs to the protein prenyltransferase subunit beta family.,similarity:Contains 5 PFTB repeats.,subunit:Heterodimer of an alpha



and a beta subunit.,


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from RAW264.7 and NIH/3T3 cells, using FNTB Antibody. The lane on the right is blocked with the synthesized peptide.