

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



MRP-L16 Polyclonal Antibody

| Catalog No | YP-Ab-03979 |
|--------------------|--|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB;IHC;IF;ELISA |
| Gene Name | MRPL16 |
| Protein Name | 39S ribosomal protein L16 mitochondrial |
| Immunogen | The antiserum was produced against synthesized peptide derived from human MRPL16. AA range:171-220 |
| Specificity | MRP-L16 Polyclonal Antibody detects endogenous levels of MRP-L16 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 |
| Observed Band | 28kD |
| Cell Pathway | Mitochondrion . |
| Tissue Specificity | Lung,Pheochromocytoma,Promyelocytic leukemia, |
| Function | function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., |
| Background | Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 39S subunit protein. [provided by RefSeq, Jul 2008], |



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:Upingbio.163.com




matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images



Western Blot analysis of various cells using MRP-L16 Polyclonal Antibody diluted at 1:2000



Immunohistochemistry analysis of paraffin-embedded human heart tissue, using MRPL16 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from 293 cells, using MRPL16 Antibody. The lane on the right is blocked with the synthesized peptide.



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



