

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



MRP-S9 Polyclonal Antibody

| Catalog No | YP-Ab-04023 |
|--------------------|--|
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB;IHC;IF;ELISA |
| Gene Name | MRPS9 |
| Protein Name | 28S ribosomal protein S9 mitochondrial |
| Immunogen | The antiserum was produced against synthesized peptide derived from human MRPS9. AA range:91-140 |
| Specificity | MRP-S9 Polyclonal Antibody detects endogenous levels of MRP-S9 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | MRPS9; RPMS9; 28S ribosomal protein S9; mitochondrial; MRP-S9; S9mt |
| Observed Band | 40kD |
| Cell Pathway | Mitochondrion . |
| Tissue Specificity | Skin, |
| Function | similarity:Belongs to the ribosomal protein S9P family.,subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., |
| Background | Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein. [provided by RefSeq, Jul 2008], |



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🗷 Email:Upingbio.163.com




matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images



Western Blot analysis of various cells using MRP-S9 Polyclonal Antibody



Western blot analysis of lysates from A549 cells, using MRPS9 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).