



# O-FucT-2 Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-04048
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	IHC;IF;ELISA
<b>Gene Name</b>	POFUT2
<b>Protein Name</b>	GDP-fucose protein O-fucosyltransferase 2
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human POFUT2. AA range:361-410
<b>Specificity</b>	O-FucT-2 Polyclonal Antibody detects endogenous levels of O-FucT-2 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	POFUT2; C21orf80; FUT13; KIAA0958; GDP-fucose protein O-fucosyltransferase 2; Peptide-O-fucosyltransferase 2; O-FucT-2
<b>Observed Band</b>	
<b>Cell Pathway</b>	Endoplasmic reticulum . Golgi apparatus . Mainly located in the endoplasmic reticulum. .
<b>Tissue Specificity</b>	Isoform A is expressed in fetal liver and peripheral blood lymphocytes. Isoform B is expressed in spleen, lung, testis, bone marrow, thymus, pancreas, prostate, fetal brain, fetal liver and fetal kidney. Isoform C is expressed in brain, heart, spleen, liver, lung, stomach, testis, placenta, skin, thymus, pancreas, mammary gland, prostate, fetal brain, fetal liver and fetal heart.
<b>Function</b>	catalytic activity:Transfers an alpha-L-fucosyl residue from GDP-beta-L-fucose to the serine hydroxy group of a protein acceptor.,function:Catalyzes the reaction that attaches fucose through an O-glycosidic linkage to a conserved serine or threonine residue in thrombospondin type 1 repeats.,online information:GlycoGene database,online information:Peptide-O-fucosyltransferase 2,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 68 family.,tissue specificity:Isoform A is expressed in fetal liver and peripheral blood lymphocytes. Isoform B is expressed in spleen, lung, testis, bone marrow, thymus, pancreas, prostate, fetal brain, fetal liver and fetal kidney. Isoform C is expressed in brain, heart, spleen, liver, lung, stomach, testis, placenta, skin, thymus, pancreas, mammary gland, prostate, fetal brain, fetal liver



and fetal heart.,

**Background**

Fucose is typically found as a terminal modification of branched chain glycoconjugates, but it also exists in direct O-linkage to serine or threonine residues within cystine knot motifs in epidermal growth factor (EGF; MIM 131530)-like repeats or thrombospondin (THBS; see MIM 188060) type-1 repeats. POFUT2 is an O-fucosyltransferase that use THBS type-1 repeats as substrates (Luo et al., 2006 [PubMed 16464857]).[supplied by OMIM, Mar 2008],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Immunohistochemistry analysis of paraffin-embedded human placenta tissue, using POFUT2 Antibody. The picture on the right is blocked with the synthesized peptide.