



MMP21 (Cleaved-Ser145) rabbit pAb

Catalog No	YP-Ab-04392
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	MMP21
Protein Name	MMP21 (Cleaved-Ser145)
Immunogen	Synthesized peptide derived from human MMP21 (Cleaved-Ser145)
Specificity	This antibody detects endogenous levels of Human, Mouse MMP21 (Cleaved-Ser145, protein was cleaved amino acid sequence between 144-145)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Matrix metalloproteinase-21 (MMP-21;EC 3.4.24.-)
Observed Band	47 65kD
Cell Pathway	Secreted .
Tissue Specificity	Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.
Function	cofactor: Binds 1 zinc ion per subunit.,cofactor: Calcium.,domain: The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function: May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM: The precursor is cleaved by a furin endopeptidase.,similarity: Belongs to the peptidase M10A family.,similarity: Contains 4 hemopexin-like domains.,tissue specificity: Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,
Background	This gene encodes a member of the matrix metalloproteinase family. Proteins in this family are involved in the breakdown of extracellular matrix for both normal physiological processes, such as embryonic development, reproduction, and



tissue remodeling, and disease processes, such as asthma and tumor metastasis. The encoded protein may play an important role in embryogenesis, particularly in neuronal cells, as well as in lymphocyte development and survival. [provided by RefSeq, May 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).