



# TGF $\beta$ Mouse mAb(1F12)

|                           |  |
|---------------------------|--|
| <b>Catalog No</b>         | YP-Ab-04780  |
| <b>Isotype</b>            | IgG  |
| <b>Reactivity</b>         | Human; Mouse;Rat   |
| <b>Applications</b>       | IHC;WB   |
| <b>Gene Name</b>          | TGFB1 TGFB   |
| <b>Protein Name</b>       | Transforming growth factor beta-1, TGF- $\beta$ 1, TGF b   |
| <b>Immunogen</b>          | Synthesized peptide derived from human TGF $\beta$ 1 Polyclonal  |
| <b>Specificity</b>        | This antibody detects endogenous levels of TGF $\beta$ at Human, Mouse,Rat   |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.11% sodium azide.  |
| <b>Source</b>             | Monoclonal, Mouse  |
| <b>Purification</b>       | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.   |
| <b>Dilution</b>           | IHC-p1:50-200 ,WB 1:1000-2000  |
| <b>Concentration</b>      | 1 mg/ml  |
| <b>Purity</b>             | $\geq 90\%$  |
| <b>Storage Stability</b>  | -20°C/1 year   |
| <b>Synonyms</b>           | Caspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12]   |
| <b>Observed Band</b>      | 52kD   |
| <b>Cell Pathway</b>       | [Latency-associated peptide]: Secreted, extracellular space, extracellular matrix . ;<br>[Transforming growth factor beta-1]: Secreted .   |
| <b>Tissue Specificity</b> | Highly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158).  |
| <b>Function</b>           |  |
| <b>Background</b>         | Transforming growth factor- $\beta$ (TGF- $\beta$ ) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other growth factors. |


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).