




KKLC1 Polyclonal Antibody

Catalog No	YP-Ab-04999
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	KKLC1 CXorf61
Protein Name	Kita-kyushu lung cancer antigen 1 (KK-LC-1) (Cancer/testis antigen 83) (CT83)
Immunogen	Synthesized peptide derived from human protein . at AA range: 30-110
Specificity	KKLC1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 12kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 12kD Cell membrane ; Single-pass type II membrane protein .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 12kD Cell membrane ; Single-pass type II membrane protein . Specifically expressed in testis. Expressed by cancer cell lines.
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 12kD Cell membrane; Single-pass type II membrane protein. Specifically expressed in testis. Expressed by cancer cell lines. tissue specificity:Specifically expressed in testis. Expressed by cancer cell lines.,
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year 12kD Cell membrane; Single-pass type II membrane protein. Specifically expressed in testis. Expressed by cancer cell lines. tissue specificity:Specifically expressed in testis. Expressed by cancer cell lines., tissue specificity:Specifically expressed in testis. Expressed by cancer cell lines.,





Products Images



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night