

Website: www.upingBio.com

ITA11 Polyclonal Antibody

| Catalog No | YP-Ab-05109 |
|--------------------|--|
| lsotype | lgG |
| Reactivity | Human;Mouse |
| Applications | WB;ELISA |
| Gene Name | ITGA11 MSTP018 |
| Protein Name | Integrin alpha-11 |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 30-110 |
| Specificity | ITA11 Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 130kD |
| Cell Pathway | Membrane; Single-pass type I membrane protein. |
| Tissue Specificity | According to PubMed:10464311, highest levels of expression in uterus and heart, intermediate levels in skeletal muscle and intermediate to low levels in pancreas, kidney and placenta. According to PubMed:10486209, also found in brain, colon, lung, small intestine, stomach, testis, salivary glands, thyroid glands and prostate. Very low levels in peripheral blood lymphocytes, fetal brain and fetal liver. |
| Function | developmental stage:Strongly up-regulated in differentiating fetal muscle cells (in vitro).,domain:The integrin I-domain (insert) is a VWFA domain. Integrins with I-domains do not undergo protease cleavage.,function:Integrin alpha-11/beta-1 is a receptor for collagen.,similarity:Belongs to the integrin alpha chain family.,similarity:Contains 1 VWFA domain.,similarity:Contains 7 FG-GAP repeats.,subunit:Heterodimer of an alpha and a beta subunit. Alpha-11 associates with beta-1. Interacts with RAB21.,tissue specificity:According PubMed:10464311 highest levels in uterus and heart, intermediate levels in skeletal muscle and intermediate to low levels in pancreas, kidney and placenta. According to PubMed:10486209 also found in brain, colon, lung, small intestine, stomach, testis, salivary glands, thyroid glands and prostate. Very low levels in peripheral blood lymphocytes, fetal brain and fet |



UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 💌 Email:Upingbio.163.com

BackgroundThis gene encodes an alpha integrin. Integrins are heterodimeric integral
membrane proteins composed of an alpha chain and a beta chain. This protein
contains an I domain, is expressed in muscle tissue, dimerizes with beta 1 integrin
in vitro, and appears to bind collagen in this form. Therefore, the protein may be
involved in attaching muscle tissue to the extracellular matrix. Alternative
transcriptional splice variants have been found for this gene, but their biological
validity is not determined. [provided by RefSeq, Jul 2008],matters needing
attentionAvoid repeated freezing and thawing!Usage suggestionsThis product can be used in immunological reaction related experiments. For

Website: www.upingBio.com

Products Images

more information, please consult technical personnel.