

Website: www.upingBio.com

TIAF1 Polyclonal Antibody

Catalog No	YP-Ab-05154
lsotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	TIAF1
Protein Name	TGFB1-induced anti-apoptotic factor 1 (12 kDa TGF-beta-1-induced antiapoptotic factor)
Immunogen	Synthesized peptide derived from human protein . at AA range: 10-90
Specificity	TIAF1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	12kD
Cell Pathway	[Isoform TIAF1]: Nucleus .
Tissue Specificity	Not detectable in normal kidney and liver. Up-regulated in chronic and acute allograft rejection: expressed in the inflammatory infiltrate and in tubular epithelial cells.
Function	caution:The TIAF1 protein is coded in the 3'-UTR region of MYO18A.,function:Inhibits the cytotoxic effects of TNF-alpha and overexpressed TNF receptor adapters TRADD, FADD, and RIP. Involved in TGF-beta1 inhibition of IkappaB-alpha expression and suppression of TNF-mediated IkappaB-alpha degradation.,
Background	caution:The TIAF1 protein is coded in the 3'-UTR region of MYO18A.,function:Inhibits the cytotoxic effects of TNF-alpha and overexpressed TNF receptor adapters TRADD, FADD, and RIP. Involved in TGF-beta1 inhibition of IkappaB-alpha expression and suppression of TNF-mediated IkappaB-alpha degradation.,
matters needing attention	Avoid repeated freezing and thawing!

Thank you for your recent purchase



UpingBio technology Co.,Ltd


🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com

WebsIte: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4°over night