

Website: www.upingBio.com

WNT8A Polyclonal Antibody

YP-Ab-05163 IgG Human;Mouse WB;ELISA WNT8A WNT8D Protein Wnt-8a (Protein Wnt-8d) Synthesized peptide derived from human protein . at AA range: 90-170 WNT8A Polyclonal Antibody detects endogenous levels of protein. Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Polyclonal, Rabbit,IgG
Human;Mouse WB;ELISA WNT8A WNT8D Protein Wnt-8a (Protein Wnt-8d) Synthesized peptide derived from human protein . at AA range: 90-170 WNT8A Polyclonal Antibody detects endogenous levels of protein. Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Polyclonal, Rabbit,IgG
WB;ELISA WNT8A WNT8D Protein Wnt-8a (Protein Wnt-8d) Synthesized peptide derived from human protein . at AA range: 90-170 WNT8A Polyclonal Antibody detects endogenous levels of protein. Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Polyclonal, Rabbit,IgG
WNT8A WNT8D Protein Wnt-8a (Protein Wnt-8d) Synthesized peptide derived from human protein . at AA range: 90-170 WNT8A Polyclonal Antibody detects endogenous levels of protein. Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Polyclonal, Rabbit,IgG
Protein Wnt-8a (Protein Wnt-8d) Synthesized peptide derived from human protein . at AA range: 90-170 WNT8A Polyclonal Antibody detects endogenous levels of protein. Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Polyclonal, Rabbit,IgG
Synthesized peptide derived from human protein . at AA range: 90-170 WNT8A Polyclonal Antibody detects endogenous levels of protein. Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Polyclonal, Rabbit,IgG
WNT8A Polyclonal Antibody detects endogenous levels of protein. Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Polyclonal, Rabbit,IgG
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Polyclonal, Rabbit,IgG
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by
affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000 ELISA 1:5000-20000
1 mg/ml
≥90%
-20°C/1 year
38kD
Secreted, extracellular space, extracellular matrix . Secreted .
function:Ligand for members of the frizzled family of seven transmembrane receptors. May play an important role in the development and differentiation of certain forebrain structures, notably the hippocampus.,similarity:Belongs to the Wnt family.,
The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family and may be implicated in development of early embryos as well as germ cell tumors. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2014],
Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd


🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com

WebsIte: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4°over night