



# ATS14 Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-05288
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	ADAMTS14
<b>Protein Name</b>	A disintegrin and metalloproteinase with thrombospondin motifs 14 (ADAM-TS 14) (ADAM-TS14) (ADAMTS-14) (EC 3.4.24.-)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 190-270
<b>Specificity</b>	ATS14 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	134kD
<b>Cell Pathway</b>	Secreted, extracellular space, extracellular matrix .
<b>Tissue Specificity</b>	Expressed in retina and at low levels in brain, lung and placenta (PubMed:11779638). High expression in fetal tissues (PubMed:11867212).
<b>Function</b>	domain:The spacer domain and the TSP type-1 domains are important for a tight interaction with the extracellular matrix.,function:Has a aminoprocollagen type 1 activity processing activity in the absence of ADAMTS2. Seems to be synthesized as a latent enzyme that requires activation to display aminoprocollagen peptidase activity.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Contains 1 disintegrin domain.,similarity:Contains 1 peptidase M12B domain.,similarity:Contains 1 PLAC domain.,similarity:Contains 4 TSP type-1 domains.,tissue specificity:Expressed in retina and at low levels in brain, lung and placenta. High expression in fetal tissues.,
<b>Background</b>	This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motif) protein family. Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The encoded preproprotein is



proteolytically processed to generate the mature enzyme. This enzyme cleaves amino-terminal propeptides from type I procollagen, a necessary step in the formation of collagen fibers. Mutations in this gene may be associated with osteoarthritis in human patients. [provided by RefSeq, May 2016],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**