

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



APOC4 Polyclonal Antibody

Catalog No YP-Ab-05326 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name APOC4 Protein Name Apolipoprotein C-IV (Apo-CIV) (ApoC-IV) (Apolipoprotein C4) Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity APOC4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/I year Synonyms Secreted. Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.similarity:Belongs to the apolipoprotein C4 family, tissue specificity:Expressed by the liver and secreted in plasma. Background This gene encodes a lipid-bind		
Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name APOC4 Protein Name Apolipoprotein C-IV (Apo-CIV) (ApoC-IV) (Apolipoprotein C4) Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity APOC4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family, tissue specificity:Expressed by the liver and secreted in plasma. Background In the protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with the related processing and the present in a cluster with other related processing and the present plasman and proprotein ct-I (APOC2) gene. [provided by RefSeq, Mar 2011],	Catalog No	YP-Ab-05326
Applications WB;ELISA Gene Name APOC4 Protein Name Apolipoprotein C-IV (Apo-CIV) (ApoC-IV) (Apolipoprotein C4) Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity APOC4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism ,similarity:Belongs to the apolipoprotein C4 family, tissue specificity:Expressed by the liver and secreted in plasma. Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein cell (APOC2) gene. [provided by RefSeq, Mar 2011],	Isotype	IgG
Gene Name APOC4 Protein Name Apolipoprotein C-IV (Apo-CIV) (ApoC-IV) (Apolipoprotein C4) Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity APOC4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism, similarity:Belongs to the apolipoprotein C4 family, tissue specificity:Expressed by the liver and secreted in plasma. Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene may influence circulating lipid levels and may be associated with orionary artery disease risk. This gene is present in a cluster with other related apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011],	Reactivity	Human;Rat;Mouse;
Protein Name Apolipoprotein C-IV (Apo-CIV) (ApoC-IV) (Apolipoprotein C4) Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity APOC4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism., similarity:Belongs to the apolipoprotein C4 family., tissue specificity:Expressed by the liver and secreted in plasma. Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011],	Applications	WB;ELISA
Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity APOC4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism., similarity:Belongs to the apolipoprotein C4 family., tissue specificity:Expressed by the liver and secreted in plasma. Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated	Gene Name	APOC4
Specificity APOC4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family., tissue specificly:Expressed by the liver and secreted in plasma., Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Protein Name	Apolipoprotein C-IV (Apo-CIV) (ApoC-IV) (Apolipoprotein C4)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function: May participate in lipoprotein metabolism., simillarity: Belongs to the apolipoprotein C4 familly., tissue specificity: Expressed by the liver and secreted in plasma., Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C1 (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human protein . at AA range: 50-130
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism., similarity:Belongs to the apolipoprotein C4 family., tissue specificity:Expressed by the liver and secreted in plasma. Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Specificity	APOC4 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 familly.,tissue specificity:Expressed by the liver and secreted in plasma., Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in plasma. Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apollipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], Matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in plasma. Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in plasma., Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Synonyms Observed Band 13kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in plasma., Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Observed Band Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in plasma., Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in plasma., Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma. Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in plasma., Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Expressed by the liver and secreted in plasma. function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in plasma., Background This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Observed Band	13kD
Function function:May participate in lipoprotein metabolism.,similarity:Belongs to the apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in plasma., This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Cell Pathway	Secreted.
apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in plasma., This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Expressed by the liver and secreted in plasma.
family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream apolipoprotein C-II (APOC2) gene. [provided by RefSeq, Mar 2011], matters needing Avoid repeated freezing and thawing!	Function	apolipoprotein C4 family.,tissue specificity:Expressed by the liver and secreted in
	Background	family. The protein is thought to play a role in lipid metabolism. Polymorphisms in this gene may influence circulating lipid levels and may be associated with coronary artery disease risk. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring downstream
	matters needing attention	Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd


📞 Tel: 400-999-8863 🗷 Emall:Upingbio.163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4° over night