

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



CTSRB Polyclonal Antibody

Catalog No	YP-Ab-05441
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	CATSPERB C14orf161
Protein Name	Cation channel sperm-associated protein subunit beta (CatSper-beta)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	CTSRB Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
•	
Synonyms	
	122kD
Synonyms	122kD Cell projection, cilium, flagellum membrane; Single-pass membrane protein. Predominantly located in the principal piece of the sperm tail.
Synonyms Observed Band	Cell projection, cilium, flagellum membrane ; Single-pass membrane protein .
Synonyms Observed Band Cell Pathway	Cell projection, cilium, flagellum membrane ; Single-pass membrane protein . Predominantly located in the principal piece of the sperm tail
Synonyms Observed Band Cell Pathway Tissue Specificity	Cell projection, cilium, flagellum membrane; Single-pass membrane protein. Predominantly located in the principal piece of the sperm tail Epithelium,Placenta,Testis, function:Probably involved in sperm cell hyperactivation via its association with CATSPER1. Sperm cell hyperactivation is needed for sperm motility which is essential late in the preparation of sperm for fertilization.,subunit:Component of a
Synonyms Observed Band Cell Pathway Tissue Specificity Function	Cell projection, cilium, flagellum membrane; Single-pass membrane protein. Predominantly located in the principal piece of the sperm tail Epithelium, Placenta, Testis, function: Probably involved in sperm cell hyperactivation via its association with CATSPER1. Sperm cell hyperactivation is needed for sperm motility which is essential late in the preparation of sperm for fertilization., subunit: Component of a complex, at least composed of CATSPER1, HSPA1B and CATSPERB., function: Probably involved in sperm cell hyperactivation via its association with CATSPER1. Sperm cell hyperactivation is needed for sperm motility which is essential late in the preparation of sperm for fertilization., subunit: Component of a
Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Cell projection, cilium, flagellum membrane; Single-pass membrane protein. Predominantly located in the principal piece of the sperm tail Epithelium, Placenta, Testis, function: Probably involved in sperm cell hyperactivation via its association with CATSPER1. Sperm cell hyperactivation is needed for sperm motility which is essential late in the preparation of sperm for fertilization., subunit: Component of a complex, at least composed of CATSPER1, HSPA1B and CATSPERB., function: Probably involved in sperm cell hyperactivation via its association with CATSPER1. Sperm cell hyperactivation is needed for sperm motility which is essential late in the preparation of sperm for fertilization., subunit: Component of a complex, at least composed of CATSPER1, HSPA1B and CATSPERB.,



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Products Images