

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com





## **CLCN2** Polyclonal Antibody

Catalog No	YP-Ab-05459	
Isotype	IgG	
Reactivity	Human;Mouse;Rat	
Applications	WB;ELISA	
Gene Name	CLCN2	
Protein Name	Chloride channel protein 2 (CIC-2)	
Immunogen	Synthesized peptide derived from part region of human protein	
Specificity	CLCN2 Polyclonal Antibody detects endogenous levels of protein.	
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Source	Polyclonal, Rabbit,IgG	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.	
Dilution	WB 1:500-2000 ELISA 1:5000-20000	
Concentration	1 mg/ml	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Synonyms		
Observed Band	98kD	
Cell Pathway	Cell membrane ; Multi-pass membrane protein.	
Tissue Specificity	Ubiquitously expressed. Moderately expressed in aortic and coronary vascular smooth muscle cells and expressed at a low level in aortic endothelial cells. Expressed in the adrenal gland, predominantly in the zona glomerulosa (PubMed:29403011).	
Function	disease:Defects in CLCN2 are a cause of juvenile absence epilepsy (JAE) [MIM:607631]. JAE is a subtype of idiopathic generalized epilepsy (IGE) characterized by onset occurring around puberty, absence seizures, generalized tonic-clonic seizures (GTCS), GTCS on awakening and myoclonic seizures.,disease:Defects in CLCN2 are the cause of childhood absence epilepsy type 3 (ECA3) [MIM:607682]. ECA3 is a subtype of idiopathic generalized epilepsy (IGE) characterized by onset at age 6-7 years, frequent absence seizures (several per day) and bilateral, synchronous, symmetric 3 Hz spike waves on EEG. During adolescence, tonic-clonic and myoclonic seizures develop.,disease:Defects in CLCN2 may be the cause of epilepsy with grand mal seizures on awakening (EGMA) [MIM:607628]. EGMA is a subtype of idiopathic generalized epilepsy (IGE) characterized by generalized tonic-clonic seizures (GTCS) occurri	



## UpingBio technology Co.,Ltd

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Background	chloride voltage-gated channel 2(CLCN2) Homo sapiens This gene encodes a voltage-gated chloride channel. The encoded protein is a transmembrane protein that maintains chloride ion homeostasis in various cells. Defects in this gene may be a cause of certain epilepsies. Four transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2012],	
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	

Products Images