



DHB8 Polyclonal Antibody

Catalog No	YP-Ab-05581
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	HSD17B8 FABGL HKE6 RING2
Protein Name	Estradiol 17-beta-dehydrogenase 8 (EC 1.1.1.62) (17-beta-hydroxysteroid dehydrogenase 8) (17-beta-HSD 8) (3-oxoacyl-[acyl-carrier-protein] reductase) (EC 1.1.1.-) (Protein Ke6) (Ke-6) (Really interest
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DHB8 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	28kD
Cell Pathway	Mitochondrion matrix .
Tissue Specificity	Widely expressed, particularly abundant in prostate, placenta and kidney (PubMed:17978863). Expressed at protein level in various tissues like brain, cerebellum, heart, lung, kidney, ovary, testis, adrenals and prostate (PubMed:30508570).
Function	catalytic activity:Estradiol-17-beta + NAD(P)(+) = estrone + NAD(P)H.,catalytic activity:Testosterone + NAD(+) = androst-4-ene-3,17-dione + NADH.,function:Uses estradiol as its preferred substrate.,pathway:Steroid biosynthesis; estrogen biosynthesis.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,tissue specificity:High expression in the liver and pancreas, lower in the skeletal muscle and kidney.,
Background	hydroxysteroid 17-beta dehydrogenase 8(HSD17B8) Homo sapiens In mice, the Ke6 protein is a 17-beta-hydroxysteroid dehydrogenase that can regulate the concentration of biologically active estrogens and androgens. It is preferentially an oxidative enzyme and inactivates estradiol, testosterone, and dihydrotestosterone. However, the enzyme has some reductive activity and can synthesize estradiol from estrone. The protein encoded by this gene is similar to



Ke6 and is a member of the short-chain dehydrogenase superfamily. An alternatively spliced transcript of this gene has been detected, but the full-length nature of this variant has not been determined. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night