

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



AGO3 Polyclonal Antibody

Catalog No	YP-Ab-05582	
Isotype	IgG	
Reactivity	Human;Mouse	
Applications	WB;ELISA	
Gene Name	EIF2C3 AGO3	
Protein Name	Protein argonaute-3 (Argonaute3) (hAgo3) (Eukaryotic translation initiation factor 2C 3) (eIF-2C 3) (eIF2C 3)	
Immunogen	Synthesized peptide derived from part region of human protein	
Specificity	AGO3 Polyclonal Antibody detects endogenous levels of protein.	
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Source	Polyclonal, Rabbit,IgG	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.	
Dilution	WB 1:500-2000 ELISA 1:5000-20000	
Concentration	1 mg/ml	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Synonyms		
Observed Band	94kD	
Cell Pathway	Cytoplasm, P-body .	
Tissue Specificity	Placenta,Testis,	
Function	function:Required for RNA-mediated gene silencing (RNAi). Binds to short RNAs such as microRNAs (miRNAs) and represses the translation of mRNAs which are complementary to them. Lacks endonuclease activity and does not appear to cleave target mRNAs.,similarity:Belongs to the argonaute family. Ago subfamily.,similarity:Contains 1 PAZ domain.,similarity:Contains 1 Piwi domain.,subunit:Interacts with EIF4B, IMP8, PRMT5 and TNRC6B.,	
Background	This gene encodes a member of the Argonaute family of proteins which play a role in RNA interference. The encoded protein is highly basic, contains a PAZ domain and a PIWI domain, and may play a role in short-interfering-RNA-mediated gene silencing. This gene is located on chromosome 1 in a tandem cluster of closely related family members including	



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



matters	needing
attentio	

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images