

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com





## TF2LY Polyclonal Antibody

Catalog No	YP-Ab-05671
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	TGIF2LY TGIFLY
Protein Name	Homeobox protein TGIF2LY (TGF-beta-induced transcription factor 2-like protein) (TGFB-induced factor 2-like protein, Y-linked) (TGIF-like on the Y)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TF2LY Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	20kD
Cell Pathway	Nucleus .
Tissue Specificity	Specifically expressed in adult testis.
Function	function:May have a transcription role in testis. May act as a competitor/regulator of TGIF2LX.,similarity:Belongs to the TALE/TGIF homeobox family.,similarity:Contains 1 homeobox DNA-binding domain.,tissue specificity:Specifically expressed in adult testis.,
Background	This gene encodes a member of the TALE/TGIF homeobox family of transcription factors. This gene lies within the male specific region of chromosome Y, in a block of sequence that is thought to be the result of a large X-to-Y transposition. The C-terminus of this protein is divergent from that of its chromosome X homolog (TGIF2LX), suggesting that this protein may act as a regulator of TGIF2LX. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!



## UpingBio technology Co.,Ltd


📞 Tel: 400-999-8863 🗷 Emall:Upingbio.163.com



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night