

Website: www.upingBio.com

RM27 Polyclonal Antibody

Catalog No	YP-Ab-05735
lsotype	lgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	MRPL27 HSPC250
Protein Name	39S ribosomal protein L27, mitochondrial (L27mt) (MRP-L27)
Immunogen	Synthesized peptide derived from human protein . at AA range: 40-120
Specificity	RM27 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	16kD
Cell Pathway	Mitochondrion .
Tissue Specificity	Brain,Lung,Umbilical cord blood,
Function	similarity:Belongs to the ribosomal protein L27P family.,
Background	Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in provide a comparison in biochemical properties, which provents approximate approximate and provents approximate the proteins comprising the mitoribosome differ greatly
	in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 39S subunit protein. [provided by RefSeq, Jul 2008],



UpingBio technology Co.,Ltd


🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com

@ Website: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4°over night