

Website: www.upingBio.com

MPP8 Polyclonal Antibody


| Catalog No | YP-Ab-05749 |
|---------------------------|--|
| lsotype | lgG |
| Reactivity | Human;Rat;Mouse |
| Applications | WB;ELISA |
| Gene Name | MPHOSPH8 MPP8 |
| Protein Name | M-phase phosphoprotein 8 (Two hybrid-associated protein 3 with RanBPM) (Twa3) |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 290-370 |
| Specificity | MPP8 Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 94kD |
| Cell Pathway | Nucleus . Chromosome . Detected on heterochromatin (PubMed:20871592, PubMed:26022416). Dissociates from chromatin during interphase and early mitosis (PubMed:23416073). Detected on nucleosomes (PubMed:20871592) |
| Tissue Specificity | Epithelium,Liver,Lymph node,Lymphoblast,Pancreas,PNS, |
| Function | PTM:Phosphorylated in M (mitotic) phase.,similarity:Contains 1 chromo domain.,similarity:Contains 4 ANK repeats., |
| Background | PTM:Phosphorylated in M (mitotic) phase.,similarity:Contains 1 chromo domain.,similarity:Contains 4 ANK repeats., |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| | |



🔇 Tel: 400-999-8863 📼 Emall:Upingbio.163.com

Website: www.upingBio.com

Products Images



Western Blot analysis of Hela lysis, using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000