



# NAV3 Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-05842
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	IHC;IF
<b>Gene Name</b>	NAV3 KIAA0938 POMFIL1 STEERIN3
<b>Protein Name</b>	Neuron navigator 3 (Pore membrane and/or filament-interacting-like protein 1) (Steerin-3) (Unc-53 homolog 3) (unc53H3)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 430-510
<b>Specificity</b>	NAV3 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	IHC-p 1:50-300. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	262kD
<b>Cell Pathway</b>	Nucleus outer membrane .
<b>Tissue Specificity</b>	Highly expressed in brain. Expressed at low levels in heart and placenta. Present in activated T-cells but not in resting T-cells (at protein level). Down-regulated in primary neuroblastoma.
<b>Function</b>	developmental stage:Expressed in fetal brain.,disease:Defects in NAV3 may be a cause of mycosis fungoides [MIM:254400]; a malignant T-cell lymphoma of the skin.,disease:Defects in NAV3 may be a cause of Sezary syndrome, a malignant T-cell lymphoma of the skin.,function:May regulate IL2 production by T-cells. May be involved in neuron regeneration.,similarity:Belongs to the Nav/unc-53 family.,similarity:Contains 1 CH (calponin-homology) domain.,tissue specificity:Highly expressed in brain. Expressed at low levels in heart and placenta. Present in activated T-cells but not in resting T-cells (at protein level). Down-regulated in primary neuroblastoma.,
<b>Background</b>	This gene belongs to the neuron navigator family and is expressed predominantly in the nervous system. The encoded protein contains coiled-coil domains and a conserved AAA domain characteristic for ATPases associated with a variety of cellular activities. This gene is similar to unc-53, a Caenorhabditis elegans gene involved in axon guidance. Multiple alternatively spliced transcript



variants for this gene have been described but only one has had its full-length nature determined. [provided by RefSeq, Jul 2008],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Immunohistochemical analysis of paraffin-embedded human Small intestinal stromal tumor. 1, Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200(4° overnight.3,Secondary antibody was diluted at 1:200(room temperature, 45min).