



PDP1 Polyclonal Antibody

Catalog No	YP-Ab-06000
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	PDP1 PDP PPM2C
Protein Name	[Pyruvate dehydrogenase [acetyl-transferring]]-phosphatase 1, mitochondrial (PDP 1) (EC 3.1.3.43) (Protein phosphatase 2C) (Pyruvate dehydrogenase phosphatase catalytic subunit 1) (PDPC 1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 180-260
Specificity	PDP1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	59kD
Cell Pathway	Mitochondrion matrix .
Tissue Specificity	Adrenal gland,Skin,Testis,
Function	catalytic activity:[Pyruvate dehydrogenase (acetyl-transferring)] phosphate + H(2)O = [pyruvate dehydrogenase (acetyl-transferring)] + phosphate.,cofactor:binds 2 magnesium ions per subunit.,disease:Defects in PDP1 are the cause of pyruvate dehydrogenase phosphatase deficiency (PDP deficiency) [MIM:608782]. PDP deficiency results in lactic acidosis leading to neurological dysfunction.,function:Catalyzes the dephosphorylation and concomitant reactivation of the alpha subunit of the E1 component of the pyruvate dehydrogenase complex.,similarity:Belongs to the PP2C family.,subunit:Heterodimer of a catalytic (PDP1) and a regulatory (PDPR) subunit.,
Background	Pyruvate dehydrogenase (E1) is one of the three components (E1, E2, and E3) of the large pyruvate dehydrogenase complex. Pyruvate dehydrogenase kinases catalyze phosphorylation of serine residues of E1 to inactivate the E1 component and inhibit the complex. Pyruvate dehydrogenase phosphatases catalyze the



dephosphorylation and activation of the E1 component to reverse the effects of pyruvate dehydrogenase kinases. Pyruvate dehydrogenase phosphatase is a heterodimer consisting of catalytic and regulatory subunits. Two catalytic subunits have been reported; one is predominantly expressed in skeletal muscle and another one is much more abundant in the liver. The catalytic subunit, encoded by this gene, is the former, and belongs to the protein phosphatase 2C (PP2C) superfamily. Along with the pyruvate dehydrogenase complex and pyruvate dehydrogenase kinases, this enzy

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images