

Website: www.upingBio.com

SSA27 Polyclonal Antibody

| Catalog No | YP-Ab-06185 |
|--|---|
| lsotype | lgG |
| Reactivity | Human;Mouse |
| Applications | WB;ELISA |
| Gene Name | SSSCA1 |
| Protein Name | Sjoegren syndrome/scleroderma autoantigen 1 (Autoantigen p27) |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | SSA27 Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| | |
| Concentration | 1 mg/ml |
| Concentration Purity | 1 mg/ml ≥90% |
| | - |
| Purity | ≥90% |
| Purity Storage Stability | ≥90% |
| Purity Storage Stability Synonyms | ≥90% -20°C/1 year |
| Purity Storage Stability Synonyms Observed Band | ≥90% -20°C/1 year |
| Purity Storage Stability Synonyms Observed Band Cell Pathway | ≥90% -20°C/1 year 21kD |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity | ≥90% -20°C/1 year 21kD Ovary, function:Might play a role in mitosis. Antigenic molecule. Could be a centromere-associated protein. May induce anti-centromere |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function | ≥90% -20°C/1 year 21kD Ovary, function:Might play a role in mitosis. Antigenic molecule. Could be a centromere-associated protein. May induce anti-centromere antibodies.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., This antigen is recognized by a subset of anti-centromere antibodies from patients with scleroderma and/or Sjogren's syndrome. Subcellular |



🕲 Tel: 400-999-8863 📼 Emall:Upingbio.163.com

Ø Website: www.upingBio.com

Products Images

