

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



SC5A5 Polyclonal Antibody

Catalog No	YP-Ab-06211	
Isotype	IgG	
Reactivity	Human;Rat;Mouse;	
Applications	WB;ELISA	
Gene Name	SLC5A5 NIS	
Protein Name	Sodium/iodide cotransporter (Na(+)/I(-) cotransporter) (Sodium-iodide symporter) (Na(+)/I(-) symporter) (Solute carrier family 5 member 5)	
Immunogen	Synthesized peptide derived from part region of human protein	
Specificity	SC5A5 Polyclonal Antibody detects endogenous levels of protein.	
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Source	Polyclonal, Rabbit,IgG	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.	
Dilution	WB 1:500-2000 ELISA 1:5000-20000	
Concentration	1 mg/ml	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Synonyms		
Observed Band	70kD	
Cell Pathway	Membrane; Multi-pass membrane protein.	
Tissue Specificity	Expression is primarily in thyroid tissue, but also to a lower extent in mammary gland and ovary. Expression is reduced in tumors.	
Function	disease:Defects in SLC5A5 are the cause of congenital hypothyroidism due to dyshormonogenesis type 1 (CHDH1) [MIM:274400]. CHDH1 is characterized by an inability of the thyroid to maintain a concentration difference of readily exchangeable iodine between the plasma and the thyroid gland, leading to congenital hypothyroidism.,function:Mediates iodide uptake in the thyroid gland.,similarity:Belongs to the sodium:solute symporter (SSF) (TC 2.A.21) family.,tissue specificity:Expression is primarily in thyroid tissue, but also to a lower extent in mammary gland and ovary. Expression is reduced in tumors.,	
Background	This gene encodes a member of the sodium glucose cotransporter family. The encoded protein is responsible for the uptake of iodine in tissues such as the thyroid and lactating breast tissue. The iodine taken up by the thyroid is incorporated into the metabolic regulators triiodothyronine (T3) and tetraiodothyronine (T4). Mutations in this gene are associated with thyroid dyshormonogenesis 1.[provided by RefSeq, Sep 2009],	



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



matters needing	
attention	

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

P	roducts Images