

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com





UBP29 Polyclonal Antibody

Catalog No	YP-Ab-06339
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	USP29
Protein Name	Ubiquitin carboxyl-terminal hydrolase 29 (EC 3.4.19.12) (Deubiquitinating enzyme 29) (Ubiquitin thioesterase 29) (Ubiquitin-specific-processing protease 29)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	UBP29 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Concentration	i mg/mi
Purity	≥90%
	<u> </u>
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 101kD Cytoplasm, perinuclear region . Localizes to perinuclear region in response to
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 101kD Cytoplasm, perinuclear region . Localizes to perinuclear region in response to
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 101kD Cytoplasm, perinuclear region . Localizes to perinuclear region in response to herpes simplex virus-1 (HSV-1) infection catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 101kD Cytoplasm, perinuclear region . Localizes to perinuclear region in response to herpes simplex virus-1 (HSV-1) infection catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family., catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year 101kD Cytoplasm, perinuclear region . Localizes to perinuclear region in response to herpes simplex virus-1 (HSV-1) infection catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family., catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,



UpingBio technology Co.,Ltd

🕻 Tel: 400-999-8863 🗷 Email:Upingbio.163.com



Products Images	