

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com





UBP3 Polyclonal Antibody

Catalog No	YP-Ab-06340
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	USP3
Protein Name	Ubiquitin carboxyl-terminal hydrolase 3 (EC 3.4.19.12) (Deubiquitinating enzyme 3) (Ubiquitin thioesterase 3) (Ubiquitin-specific-processing protease 3)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	UBP3 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year 57kD
Synonyms	ř
Synonyms Observed Band	57kD
Synonyms Observed Band Cell Pathway	57kD Nucleus . Localizes preferentially with monoubiquitinated H2A to chromatin.
Synonyms Observed Band Cell Pathway Tissue Specificity	Nucleus . Localizes preferentially with monoubiquitinated H2A to chromatin. Expressed in all tissues examined, with strongest expression in pancreas. catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 1 UBP-type zinc finger.,tissue specificity:Expressed in all tissues examined, with
Synonyms Observed Band Cell Pathway Tissue Specificity Function	Nucleus . Localizes preferentially with monoubiquitinated H2A to chromatin. Expressed in all tissues examined, with strongest expression in pancreas. catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., similarity: Contains 1 UBP-type zinc finger., tissue specificity: Expressed in all tissues examined, with strongest expression in pancreas., catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., similarity: Contains 1 UBP-type zinc finger., tissue specificity: Expressed in all tissues examined, with
Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Nucleus . Localizes preferentially with monoubiquitinated H2A to chromatin. Expressed in all tissues examined, with strongest expression in pancreas. catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 1 UBP-type zinc finger.,tissue specificity:Expressed in all tissues examined, with strongest expression in pancreas., catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 1 UBP-type zinc finger.,tissue specificity:Expressed in all tissues examined, with strongest expression in pancreas.,



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Products Images