

Website: www.upingBio.com

## ZBT10 Polyclonal Antibody

Catalog No	YP-Ab-06403
Isotype	lgG
Reactivity	Human;Rat
Applications	WB;ELISA
Gene Name	ZBTB10 RINZF RINZFC
Protein Name	Zinc finger and BTB domain-containing protein 10 (Zinc finger protein RIN ZF)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	ZBT10 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 95kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 95kD Nucleus .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 95kD Nucleus . Brain,PCR rescued clones,Prostate,Testis, function:May be involved in transcriptional regulation.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 BTB (POZ)
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 95kD Nucleus . Brain,PCR rescued clones,Prostate,Testis, function:May be involved in transcriptional regulation.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 2 C2H2-type zinc fingers., function:May be involved in transcriptional regulation.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 2 C2H2-type zinc fingers.,



🔇 Tel: 400-999-8863 📼 Emall:Upingbio.163.com

Ø Website: www.upingBio.com

## **Products Images**