

Website: www.upingBio.com

ZBTB2 Polyclonal Antibody

Catalog No	YP-Ab-06404
lsotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	ZBTB2 KIAA1483 ZNF437
Protein Name	Zinc finger and BTB domain-containing protein 2
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	ZBTB2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 56kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 56kD Nucleus .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 56kD Nucleus . Brain,Melanoma,Stomach, function:May be involved in transcriptional regulation.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 BTB (POZ)
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 56kD Nucleus . Brain,Melanoma,Stomach, function:May be involved in transcriptional regulation.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 4 C2H2-type zinc fingers., function:May be involved in transcriptional regulation.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 4 C2H2-type zinc fingers.,



🔇 Tel: 400-999-8863 📼 Emall:Upingbio.163.com

Ø Website: www.upingBio.com

Products Images