

Tel: 400-999-8863
 ■ Email:Upingbio.163.com



ZN335 Polyclonal Antibody

Catalog NoYP-Ab-06413IsotypeIgGReactivityHuman;Rat;MouseApplicationsWB;ELISAGene NameZNF335Protein NameZinc finger protein 335 (NRC-interacting factor 1) (NIF-1)ImmunogenSynthesized peptide derived from part region of human proteinSpecificityZN335 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgG	
Reactivity Human;Rat;Mouse Applications WB;ELISA Gene Name ZNF335 Protein Name Zinc finger protein 335 (NRC-interacting factor 1) (NIF-1) Immunogen Synthesized peptide derived from part region of human protein Specificity ZN335 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Applications WB;ELISA Gene Name ZNF335 Protein Name Zinc finger protein 335 (NRC-interacting factor 1) (NIF-1) Immunogen Synthesized peptide derived from part region of human protein Specificity ZN335 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Gene Name ZNF335 Protein Name Zinc finger protein 335 (NRC-interacting factor 1) (NIF-1) Immunogen Synthesized peptide derived from part region of human protein Specificity ZN335 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Protein Name Zinc finger protein 335 (NRC-interacting factor 1) (NIF-1) Immunogen Synthesized peptide derived from part region of human protein ZN335 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Immunogen Synthesized peptide derived from part region of human protein Specificity ZN335 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Specificity ZN335 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
D. I. J. D. II. (1. C.	
Source Polyclonal, Rabbit,IgG	
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.	
Dilution WB 1:500-2000 ELISA 1:5000-20000	
Concentration 1 mg/ml	
Purity ≥90%	
Storage Stability -20°C/1 year	
Synonyms	
Observed Band 147kD	
Cell Pathway Nucleus .	
Tissue Specificity Ubiquitously expressed.	
function: May regulate transcriptional activation through NCOA6.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 13 C2H2-type zinc fingers.,subunit:Interacts wi NCOA6, but not with ligand-bound nuclear hormone receptors, including THRA, RARA, RXRA, GCCR and PPARA.,tissue specificity:Relatively hig expression in the skeletal muscle, thymus, placenta and blood. Moderate expression in the colon, kidney and lung. Low expression in the small into heart, liver and brain.,	ESR1, gh
zinc finger protein 335(ZNF335) Homo sapiens The protein encoded gene enhances transcriptional activation by ligand-bound nuclear hormor receptors. However, it does this not by direct interaction with the receptor direct interaction with the nuclear hormone receptor transcriptional coacti NRC. The encoded protein may function by altering local chromatin struction [provided by RefSeq, Jul 2008],	ne r, but by ivator



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



matters needing	
attention	

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

P	roducts Images