

Website: www.upingBio.com

CDK19 Polyclonal Antibody

Catalog No	YP-Ab-06440
Isotype	lgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	CDK19 CDC2L6 CDK11 KIAA1028
Protein Name	Cyclin-dependent kinase 19 (EC 2.7.11.22) (CDC2-related protein kinase 6) (Cell division cycle 2-like protein kinase 6) (Cell division protein kinase 19) (Cyclin-dependent kinase 11) (Death-preventing
Immunogen	Synthesized peptide derived from human protein . at AA range: 30-110
Specificity	CDK19 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	55kD
Cell Pathway	Cytoplasm . Cytoplasm, perinuclear region . Nucleus .
Tissue Specificity	Testis,
Function	protein amino acid phosphorylation, phosphorus metabolic process, phosphate metabolic process, phosphorylation,cell division,
Background	This gene encodes a protein that is one of the components of the Mediator co-activator complex. The Mediator complex is a multi-protein complex required for transcriptional activation by DNA binding transcription factors of genes transcribed by RNA polymerase II. The protein encoded by this gene is similar to cyclin-dependent kinase 8 which can also be a component of the Mediator complex. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2014],
matters needing	Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com

WebsIte: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images