

SYWM Polyclonal Antibody

| Catalog No | YP-Ab-06587 |
|------------------------------|--|
| Isotype | lgG |
| Reactivity | Human;Mouse |
| Applications | WB;ELISA |
| Gene Name | WARS2 |
| Protein Name | TryptophantRNA ligase, mitochondrial (EC 6.1.1.2) ((Mt)TrpRS) (Tryptophanyl-tRNA synthetase) (TrpRS) |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | SYWM Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 39kD |
| Cell Pathway | Mitochondrion matrix . |
| Tissue Specificity | Brain. |
| Function | catalytic activity:ATP + L-tryptophan + tRNA(Trp) = AMP + diphosphate + L-tryptophyl-tRNA(Trp).,similarity:Belongs to the class-I aminoacyl-tRNA synthetase family., |
| Background | Aminoacyl-tRNA synthetases catalyze the aminoacylation of tRNA by their cognate amino acid. Because of their central role in linking amino acids with nucleotide triplets contained in tRNAs, aminoacyl-tRNA synthetases are thought to be among the first proteins that appeared in evolution. Two forms of tryptophanyl-tRNA synthetase exist, a cytoplasmic form, named WARS, and a mitochondrial form, named WARS2. This gene encodes the mitochondrial tryptophanyl-tRNA synthetase. Two alternative transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008], |
| matters needing attention | Avoid repeated freezing and thawing! |



UpingBio technology Co.,Ltd


🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com

@ Website: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4°over night