



PLCE1 Polyclonal Antibody

Catalog No	YP-Ab-06656
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF
Gene Name	PLCE1 KIAA1516 PLCE PPLC
Protein Name	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase epsilon-1 (EC 3.1.4.11) (Pancreas-enriched phospholipase C) (Phosphoinositide phospholipase C-epsilon-1) (Phospholipase C-epsilon-1) (PLC-epsi
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	PLCE1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC-p 1:50-300. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	253kD
Cell Pathway	Cytoplasm, cytosol. Cell membrane. Golgi apparatus membrane. Cell projection, lamellipodium . Recruited to plasma membrane by activated HRAS and RAP2. Recruited to perinuclear membrane by activated RAP1A. Isoform 1 and isoform 2 associates with Golgi membranes.
Tissue Specificity	Widely expressed. Expressed in podocytes (PubMed:29058690). ; [Isoform 1]: Broadly expressed and only absent in peripheral blood leukocytes. ; [Isoform 2]: Specifically expressed in placenta, lung and spleen.
Function	catalytic activity:1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate + H(2)O = 1D-myo-inositol 1,4,5-trisphosphate + diacylglycerol.,cofactor:Calcium.,disease:Defects in PLCE1 are the cause of nephrotic syndrome type 3 (NPHS3) [MIM:610725]; also called early-onset nephrotic syndrome type 3. Nephrotic syndrome, a malfunction of the kidney glomerular filter, leads to proteinuria, hypoalbuminemia, edema. End-stage kidney disease is observed in steroid-resistant nephrotic syndrome.,domain:The Ras-associating domain 1 is degenerated and may not bind HRAS. The Ras-associating domain 2 mediates interaction with GTP-bound HRAS, RAP1A, RAP2A and RAP2B and recruitment of HRAS to the cell membrane.,domain:The Ras-GEF domain has a GEF activity towards HRAS and RAP1A. Mediates



activation of the mitogen-activated protein kinase pathway.,enzyme regulation:Activated by the heterotrimeric G-protein subunit

Background

This gene encodes a phospholipase enzyme that catalyzes the hydrolysis of phosphatidylinositol-4,5-bisphosphate to generate two second messengers: inositol 1,4,5-triphosphate (IP3) and diacylglycerol (DAG). These second messengers subsequently regulate various processes affecting cell growth, differentiation, and gene expression. This enzyme is regulated by small monomeric GTPases of the Ras and Rho families and by heterotrimeric G proteins. In addition to its phospholipase C catalytic activity, this enzyme has an N-terminal domain with guanine nucleotide exchange (GEF) activity. Mutations in this gene cause early-onset nephrotic syndrome; characterized by proteinuria, edema, and diffuse mesangial sclerosis or focal and segmental glomerulosclerosis. Alternative splicing results in multiple transcript variants encoding distinct isoforms.[provided by RefSeq, Sep 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).